



STEELHEAD FISH HATCHERY EVALUATIONS—IDAHO



T. Dean Rhine Senior Fishery Research Biologist

Randall S. Osborne Senior Fisheries Technician

Kristy A. Stevens Fisheries Biological-Aide

IDFG Report Number 99-26 March 1999

Steelhead Fish Hatchery Evaluations—Idaho

Period Covered: October 1, 1993 to September 30, 1994

Ву

T. Dean Rhine Randall S. Osborne Kristy A. Stevens

Idaho Department of Fish and Game 600 South Walnut Street P.O. Box 25 Boise, ID 83707

To

U.S. Fish and Wildlife Service Lower Snake River Compensation Plan Office 1387 S. Vinnell Way, Suite 343 Boise, ID 83709

Contract Number 14-48-0001-94500

IDFG Report Number 99-26 March 1999

TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT	1
INTRODUCTION	2
METHODS	3
IDEC LODOD Decrease Consessor Decompositories	2
IDFG LSRCP Program Success Documentation	
Hatchery Database Development Hatchery Operations Documentation	
•	
Migration Conditions Juvenile Migration Timing and Survival	44
Adult Returns	
Fisheries Contribution	
Hatchery Weirs	
Smolt-to-Adult Return Rate	
Experimentation	
Clearwater Fish Hatchery	
Cover Experiment6	
Fin Erosion Experiment6	
Hagerman National Fish Hatchery	6
Size-at-Release Experiment6	
Acclimation Experiment6	
RESULTS	7
Hatchery Database Development	7
Hatchery Operations Documentation	
Clearwater Fish Hatchery	
Brood Year 19937	
Brood Year 19948	
Hagerman National Fish Hatchery	9
Brood Year 19939	
Brood Year 19949	
Magic Valley Fish Hatchery	10
Brood Year 199310	
Brood Year 199410	
Migration Conditions	10
Migration Timing and Juvenile Survival	11
Adult Returns	
Fisheries Contribution	
Hatchery Weirs	11
Sawtooth Fish Hatchery Weir11	
East Fork Salmon River Weir11	
Smolt-to-Adult Return Rates	
Clearwater Fish Hatchery	
Hagerman National Fish Hatchery	
Magic Valley Fish Hatchery	13

Experime	entation13
	Table of Contents (Continued)
	<u>Page</u>
	vater Fish Hatchery
Hager	man National Fish Hatchery13
	ze-at-Release Experiment13 cclimation Experiment14
LITERATU	RE CITED
APPENDIC	DES
	LIST OF TABLES
Table 1.	Number of unique PIT tag interrogations of LSRCP steelhead smolts, by PIT tag file, at Lower Granite Dam (GRJ), Little Goose Dam (GOJ), Lower Monumental Dam (LMJ), and McNary Dam (MCJ) for the 1994 migration period. A total of 4,665 PIT-tagged steelhead were released from Hagerman National Fish Hatchery, Magic Valley Fish Hatchery, and Clearwater Fish Hatchery between April 5 and May 3, 1994. Median travel time is to Lower Granite Dam.17
Table 2.	Snake River mean daily in-flow (thousand cubic feet per second) at Lower Granite Dam during the Peak and Extended chinook salmon smolt migration periods, 1977-1994. The migration periods are as defined by Petrosky (1991).21
Table 3.	Estimated number of LSRCP steelhead that returned to Idaho in 1993-1994. The adult return in 1993-1994 included fish from three age classes. Steelhead were reared at Hagerman National and Magic Valley fish hatcheries. These estimates were prepared by the Idaho Department of Fish and Game's Harvest Monitoring Project and only include steelhead harvested in Idaho's sport fisheries, steelhead that returned to hatchery racks, and fish that escaped to spawn naturally. These are minimum estimates and do not include all tributary and mainstem strays or in-river pre-spawning mortalities.22
Table 4.	Steelhead smolts released from Magic Valley and Hagerman National fish hatcheries that contributed to the 1993-1994 Idaho steelhead return. The number of steelhead smolts released and the estimated number of adults that returned were compared to facility design production targets and projected adult return goals.23
Table 5.	Summary of the 1994 A-strain steelhead trout return to the Sawtooth Fish Hatchery weir. The fish return included fish of hatchery and natural origin. Hatchery aging criteria, based on length, were used to determine age ^a . ND indicates "no data" (i.e., data were not available).24

List of Tables (Continued.)

Page

- Table 6. Summary of the 1994 B-strain steelhead trout return to the East Fork Salmon River weir. The fish return included fish of hatchery and natural origin. Hatchery aging criteria, based on length, were used to determine age^a. ND indicates "no data" (i.e., data were not available).25
- Table 7. Smolt-to-adult return rates of coded-wire-tagged steelhead smolts released from Hagerman National and Magic Valley fish hatcheries. The number of adults was estimated by Ball (1997) and only include steelhead harvested in Idaho's sport fisheries, steelhead that returned to hatchery racks, and fish that escaped to spawn naturally. These are minimum estimates and do not include all tributary and mainstem strays or in-river pre-spawning mortalities.26
- Table 8. Total number^a of steelhead recovered with CWTs designating them as either large size^b or regular size^c. Recovered fish were released as smolts at the Sawtooth Fish Hatchery weir in 1991 (brood year 1990) and 1992 (brood year 1991). Recovery data were not available for the 2-ocean component of brood year 1991 (i.e., ND indicates "no data").27
- Table 9. Total number^a of steelhead recovered with CWTs designating them as either acclimated (ACC) or non-acclimated (NON-ACC). Recovered fish were released as smolts at the Sawtooth Fish Hatchery weir in 1992 (brood year 1991). ND indicates "No Data" (i.e., data were not available).27

LIST OF FIGURES

Figure 1. Number of unique interrogations of LSRCP juvenile steelhead at Lower Granite Dam, Washington, plotted with the average daily in-flow of the Snake River (at Lower Granite Dam) in 1994. A total of 4,665 PIT-tagged steelhead were released from Hagerman National, Magic Valley, and Clearwater fish hatcheries between April 5 and May 3, 1994. Thirty-seven percent (1,728) of the PIT-tagged fish were interrogated at Lower Granite Dam. Data for 16 fish fall outside of this date range and are not shown.28

LIST OF APPENDICES

Page

- Appendix A. Table 1. Release data and estimated adult returns for Clearwater Fish Hatchery summer steelhead trout, brood year 1993. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.30
- Appendix A. Table 2. Release data and estimated adult returns for Hagerman National Fish Hatchery summer steelhead trout, brood year 1993. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.34
- Appendix A. Table 3. Release data and estimated adult returns for Magic Valley Fish Hatchery summer steelhead trout, brood year 1993. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.37

List of Appendices (Continued.)

Page

- Appendix B. Table 1. Release data and estimated adult returns for Hagerman National Fish Hatchery summer steelhead trout, brood year 1989. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.42
- Appendix B. Table 2. Release data and estimated adult returns for Hagerman National Fish Hatchery summer steelhead trout, brood year 1990. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.45
- Appendix B. Table 3. Release data and estimated adult returns for Hagerman National Fish Hatchery summer steelhead trout, brood year 1991. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.48

List of Appendices (Continued.)

Page

- Appendix C. Table 1. Release data and estimated adult returns for Magic Valley Fish Hatchery summer steelhead trout, brood year 1989. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.51
- Appendix C. Table 2. Release data and estimated adult returns for Magic Valley Fish Hatchery summer steelhead trout, brood year 1990. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.54
- Appendix C. Table 3. Release data and estimated adult returns for Magic Valley Fish Hatchery summer steelhead trout, brood year 1991. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.58

ABSTRACT

This annual report summarizes activities associated with Idaho Lower Snake River Compensation Plan (LSRCP) hatcheries' activities from October 1, 1993 to September 30, 1994. Included in this report are all fall 1993 and spring 1994 adult steelhead trout *Oncorhynchus mykiss* returns and all releases of juvenile steelhead trout made within the reporting period. Information presented in this report supersedes that included in previous reports.

An estimated 7,534 LSRCP steelhead returned to Idaho during this reporting period. Magic Valley Fish Hatchery produced an estimated 4,446 adult steelhead, and Hagerman National Fish Hatchery produced an estimated 3,088 adult steelhead. Steelhead returns to Idaho remain well below the LSRCP program mitigation goal of 39,260 adults.

Adult returns to Idaho LSRCP hatchery racks included 338 A-strain steelhead to Sawtooth Fish Hatchery and 73 B-strain steelhead to the East Fork Salmon River satellite facility. Approximately 52% of the hatchery steelhead that returned to Sawtooth Fish Hatchery were males as compared to 62% of the hatchery fish that returned to the East Fork Salmon River satellite.

Coded-wire tags were used to determine smolt-to-adult return rates. Dworshak B-stock steelhead, brood year 1989, reared at Hagerman National Fish Hatchery and released into the Little Salmon River had a mean smolt-to-adult return rate of 1.22%. Large size and normal size Sawtooth A-stock steelhead smolts, brood year 1990, reared at Hagerman National Fish Hatchery as part of a size-at-release experiment had mean smolt-to-adult return rates of 0.58 and 0.46%, respectively. Dworshak B- and East Fork B-stock smolts, brood year 1989, reared at Magic Valley Fish Hatchery and released into the East Fork Salmon River had mean smolt-to-adult return rates of 0.16% and 0.22%, respectively. Pahsimeroi A-stock steelhead smolts, brood year 1989, reared at Magic Valley Fish Hatchery and released into the Little Salmon River had a mean smolt-to-adult return rate of 0.34%.

In April and May 1994, Idaho LSRCP hatcheries released 4,161,359 brood year 1993 steelhead smolts. A total of 774,977 smolts were released with coded-wire tags, and 4,665 were released with passive integrated transponder tags.

INTRODUCTION

The Water Resources Development Act of 1976 (90 Stat. 2917) authorized the Idaho Lower Snake River Compensation Plan (LSRCP) to mitigate for fish losses caused by the construction and operation of Ice Harbor, Lower Monumental, Little Goose, and Lower Granite dams on the lower Snake River. Mitigation for anadromous fishery losses included the construction and operation of fish hatchery facilities and smolt passage improvements at the lower Snake River dams. The United States Fish and Wildlife Service was authorized to administer the operation and maintenance for 12 hatcheries and 11 satellite facilities in Idaho, Oregon, and Washington.

In Idaho, the Idaho Department of Fish and Game (IDFG) operates Clearwater Fish Hatchery, McCall Fish Hatchery, Magic Valley Fish Hatchery, Sawtooth Fish Hatchery, South Fork Salmon River Trap, East Fork Salmon River Trap, and Red River, Crooked River, and Powell satellite facilities. The United States Fish and Wildlife Service operates Dworshak National Fish Hatchery and Hagerman National Fish Hatchery. Adult return goals for the entire LSRCP are 8,000 summer chinook salmon *Oncorhynchus tshawytscha*, 50,700 spring chinook salmon, 18,300 fall chinook salmon and 55,100 steelhead trout *O. mykiss* to the Snake River basin. Adult return goals for the Idaho portion of the LSRCP program call for the return of 8,000 adult summer chinook salmon, 40,432 adult spring chinook salmon, and 39,260 adult steelhead. Adult return goals for Idaho LSRCP steelhead hatcheries are as follows: Clearwater Fish Hatchery—14,000, Hagerman National Fish Hatchery—13,600, and Magic Valley Fish Hatchery—11,660.

The LSRCP includes a Hatchery Evaluation Study component to monitor and evaluate the mitigation hatchery program. The primary objective of the Hatchery Evaluation Study is to determine the best hatchery management practices for mitigation hatcheries to meet LSRCP and IDFG anadromous fisheries goals. Only if we understand the effects of hatchery operations on adult return characteristics (e.g., return rates, sex ratios, and age structure) can we prescribe effective management actions. Tasks we defined to satisfy the primary objective are divided into two categories: 1) documentation, and 2) investigation. We document hatchery practices for each brood year, or cohort, of fish and mitigation status in terms of annual adult returns. Our success at achieving LSRCP and IDFG goals can then be related to hatchery practices through the documentation tasks. Investigation tasks are manipulative experiments involving modified or alternative hatchery practices that show potential for increasing adult returns and achieving LSRCP and IDFG goals.

This report summarizes steelhead trout Hatchery Evaluation Study activities carried out from October 1, 1993 through September 30, 1994. Juvenile steelhead released from Clearwater, Hagerman National, and Magic Valley fish hatcheries during this reporting period are documented, as well as adult steelhead that returned to Idaho during the fall of 1993 and the spring of 1994 (hereafter referred to as the 1993-1994 return). Specific objectives identified in Cooperative Work Agreement 14-48-0001-94500 are covered in this report and are as follows:

Objective 1. Document the success of the IDFG-LSRCP program in meeting specific adult return goals.

Sub-objective 1.1 Develop a computerized hatchery database standardizing all necessary variables for monitoring and evaluation.

- Sub-objective 1.2 Document LSRCP fish rearing and release practices and adult returns in Idaho.
- Objective 2. Identify factors limiting hatchery success and recommend possible improvements based on existing knowledge and experimentation.
 - Sub-objective 2.1 Continue ongoing documentation and monitoring to determine the relationships between adult returns and hatchery practices, characteristics of hatchery products, and juvenile survival.
 - Sub-objective 2.2 Conduct controlled studies (short-term experiments) to determine the relationships between adult returns and hatchery practices, characteristics of hatchery products, and juvenile survival.

The results of experiments performed under Sub-objective 2.2 are printed separately from this report. Some results from those experiments, such as juvenile migration characteristics and adult return rates for experimental groups, are included in this report.

METHODS

IDFG LSRCP Program Success Documentation

To document the overall success of the program, we compared the estimated number of adult steelhead that returned above Lower Granite Dam between October 1, 1993 and September 30, 1994 to the Idaho LSRCP goal of 25,260 adult steelhead. The 39,260 fish goal was not used because 14,000 of these fish were charged to Clearwater Fish Hatchery, which only became operational in 1992. Adults from the first smolt release from Clearwater Fish Hatchery will return in 1995, 1996, and 1997. The Harvest Monitoring Project estimated the total number of returning adults and partitioned the total return between Hagerman National Fish Hatchery and Magic Valley Fish Hatchery based on coded-wire tag data. Results for Objective 1 are reported under *Results*, *Adult Returns*.

Hatchery Database Development

A team of IDFG personnel from Management, Hatcheries, Natural Resources, and the Fish Health and Coded-Wire Tag labs was organized to develop a computerized database for Idaho's anadromous hatcheries. The team will develop a detailed objective statement that describes the purpose of the database, input variables, and expected output functions. The team will ensure the database meets minimum standards for all parties. A database programmer will be hired to develop the product. Tasks for the programmer include writing computer code, developing input screens, developing summary output screens and reports, field testing the program, and preparing database documentation.

Hatchery Operations Documentation

Hatchery operations between October 1, 1993 and September 30, 1994 are documented in this report. Pertinent rearing information affecting brood years 1993 and 1994 are discussed. Additional information which occurred prior to this reporting period may be included for brood year 1993 steelhead for completeness. Information was collected from Hatchery Brood Year and Run reports, memorandums, and from verbal communications with hatchery personnel. Fish marking and tagging information was provided by IDFG's Coded-Wire Tag Lab.

Migration Conditions

Snake River flow during smolt migration is a major factor affecting the survival of Idaho's anadromous fishes. This reporting period covers adult steelhead that returned to Idaho in the fall of 1993 and the spring of 1994. Returning adults could be from brood year 1989, 1990, or 1991, depending on their age at return. Since steelhead smolts for these brood years were released the following spring, flow conditions for 1990, 1991, and 1992 are reported. Flow conditions for the 1994 emigration period are also reported. Water flow data were obtained from Fish Passage Center reports and the United States Geological Survey Internet site.

Petrosky (1991) defined two time periods that accounted for most of the chinook emigration past Lower Granite Dam. The Peak period of emigration is from April 15 to May 5 and is the time period when approximately 50% of the yearling chinook salmon reach Lower Granite Dam. The Extended period is from April 20 to May 30 and encompasses the time when most of the wild and natural yearling chinook salmon reach Lower Granite Dam. Hatchery steelhead smolts are generally released in April and emigrate during the same time period as chinook. Therefore, flows during the Extended and Peak time periods are reported.

Juvenile Migration Timing and Survival

Passive Integrated Transponder (PIT) tags were used to evaluate downstream migration. The interrogation rate of PIT-tagged juvenile salmonids at lower Snake River dams serves as a minimum survival index because: 1) an unknown (but we believe small) number of PIT-tagged fish that die in the hatchery may go undetected, although we scan the dead fish; 2) not all fish pass through detectors; 3) some PIT tags fail (approximately 2%, R. Kiefer, IDFG, personal communication) or are lost between tagging and arrival at detection sites; 4) some fish arrive while detection gear is not being operated, and 5) mortality occurs between dams.

Brood year 1993 steelhead juveniles were PIT-tagged by IDFG Fish Marking personnel. The PIT tag data were submitted to PTAGIS, a computerized PIT tag database operated by Pacific States Marine Fisheries Commission (Columbia River Basin PIT Tag Information System 1997). Interrogation rates and median travel times for specific PIT-tagged groups of steelhead were calculated after retrieving relevant interrogation data from PTAGIS. Interrogation rates were calculated for each PIT tag file (or files depending on the purpose of the tagging), by dividing the number of unique interrogations at Lower Granite, Little Goose, Lower Monumental, and McNary dams by the number of PIT-tagged fish released, multiplied by 100. Median travel times were calculated for each PIT tag file (or files) to Lower Granite Dam.

Adult Returns

Adult return goals for Clearwater Fish Hatchery, Hagerman National Fish Hatchery, and Magic Valley Fish Hatchery are 14,000, 13,600, and 11,660 adult steelhead above Lower Granite Dam, respectively. The Harvest Monitoring Project (i.e., Ball 1997) estimated the total number of LSRCP steelhead that returned to Idaho in the 1993-1994 return. This estimate included LSRCP-reared fish that were harvested in Idaho's sport fishery and LSRCP-reared fish that escaped to spawn naturally or returned to hatchery racks. Ball's (1997) estimate should be considered a minimum estimate because tributary and mainstem strays were not accounted for, nor were in-river pre-spawning mortalities. The number of steelhead smolts released and the estimated number of adults that returned were compared to facility design production targets and projected adult return goals for each facility.

Fisheries Contribution

Fish Marking personnel from IDFG tagged juvenile steelhead with coded-wire tags (CWTs) according to marking/tagging plans developed by fishery managers and research biologists. Steelhead tagged with CWTs had the left pelvic fin excised to indicate the presence of a tag. The snouts from tagged adult steelhead harvested in Idaho's sport fishery were sent to the Coded-Wire Tag Lab and processed. The Harvest Monitoring Project used these data, along with data from a statewide telephone survey, to estimate the total number of steelhead harvested in Idaho. The numbers of steelhead harvested from specific release groups were estimated by expanding codedwire tag recoveries for specific groups. See Ball (1997) for methods.

Hatchery Weirs

The numbers of steelhead that returned to the East Fork Salmon River and Sawtooth Fish Hatchery weirs were documented by hatchery personnel. The length, sex, and disposition of each fish were recorded. Fish length and strain (A or B) were used to determine age-at-return. Snouts from coded-wire-tagged steelhead were sent to the Coded-Wire Tag Lab and processed. The Harvest Monitoring Project used these data to estimate the total number of LSRCP-reared steelhead that returned to hatchery racks or escaped to spawn naturally.

Smolt-to-Adult Return Rate

The Harvest Monitoring Project estimated the total number of LSRCP-produced steelhead that returned to Idaho. For each coded-wire tag (CWT) code, we summed the estimated number of steelhead that returned to Idaho in the 1991-1992 (Ball 1994), 1992-1993 (Ball 1996), and the 1993-1994 (Ball 1997) harvest seasons. For specific groups of fish (i.e., individual CWT codes by release site), we calculated a smolt-to-adult return rate (SAR) by dividing the estimated number of adults that returned (provided by the Harvest Monitoring Project) by the number of smolts released (provided by the Coded-Wire Tag Lab), multiplied by 100.

Experimentation

Interim progress reports, printed independently of this report, document the current status of Hatchery Evaluation Study experiments. Results from some experiments, particularly adult return rates and emigration rates for experimental groups, are included in this report.

Clearwater Fish Hatchery

<u>Cover Experiment</u>—This experiment was initiated in 1992 using Dworshak B-stock steelhead from the 1992 brood. The purpose of the experiment was to test the effects of shade covers on adult return and juvenile emigration rates. Adult returns will be complete in 1997. See Rhine et al. (In Press) for details of the experimental design.

<u>Fin Erosion Experiment</u>—The purpose of the experiment was to test the effects of raceways (design and feeding method) and baffles (raceways with and without baffles) on adult steelhead return and juvenile emigration rates. Dworshak B-stock steelhead from the 1993 brood were used for the study. Contact Clearwater Fish Hatchery personnel for study design and completion report.

Hagerman National Fish Hatchery

<u>Size-at-Release Experiment</u>—The purpose of this study was to determine the optimal size (length) to rear steelhead trout juveniles at Hagerman National Fish Hatchery. The experiment was conducted for two consecutive years using A-strain steelhead from the 1990 and 1991 broods. Adult returns will be complete in 1995. See Cannamela (1992) for complete details of the experimental design.

<u>Acclimation Experiment</u>—This study compared juvenile emigration and adult return rates of steelhead that were trucked from Hagerman National Fish Hatchery and acclimated at Sawtooth Fish Hatchery for two weeks (Acclimated Group) to steelhead that were trucked from Hagerman National Fish Hatchery two weeks later and released directly into the Salmon River (Non-acclimated Group). The two groups of fish were released at the same time and at the same location. This was the third year of the study. The study was initiated in 1992 and repeated in 1993 and 1994. Pahsimeroi A-stock steelhead were used in the 1992 and 1993 study years while Sawtooth A-stock steelhead were used in the 1994 study year. Adult returns from the 1991, 1992, and 1993 broods will be complete in 1995, 1996, and 1997, respectively. Adult return data and emigration data will be analyzed by brood year. The total number of adults that return from each group will be tested using chi-square analysis (α = .05). Passive Integrated Transponder tags will be used to determine unique interrogation rates at Snake and Columbia river dams and median travel time to Lower Granite Dam for each group. Chi-square analysis will be used to test interrogation rates between groups. Travel times for the two groups will be tested for differences using the Mann-Whitney test (α = .05) (SYSTAT Inc. 1992).

RESULTS

Hatchery Database Development

A team was formed to work on the project, and a computer programmer (Nuralima Boydstun) was hired to develop the product. An objective statement was developed along with a description of the expected product. The database program was envisioned to increase efficiency, improve record keeping, and standardize hatchery operations. Each hatchery will be equipped with a version of the menu-operated program. A slightly different version, capable of compiling all of the individual hatchery records, will be operated from the Headquarters Office in Boise. Data will be transferred via modem or by diskette. The database program will document all phases of hatchery operations (e.g., trapping and incubation). It will provide the following benefits. First, the database program will serve as a working tool for hatchery personnel by providing summary output functions suitable for hatchery management and documentation tasks. Second, the database will be capable of storing many intricate, in-hatchery details that will be used for evaluation purposes. Third, the hatchery database program will serve as a standardizing mechanism for data collections among hatcheries. Finally, the program will provide easy access to hatchery data by storing data in one location and in one format.

Specification documents for the PTAGIS, Pacific States Marine Fisheries Commission, CIS, and Idaho CWT databases were obtained to standardize field names, data codes, and abbreviations. The database structure for the adult trapping and spawning components were developed. Nuralima Boydstun developed the menu input screens for the trapping portion of the database and prepared database documentation. Nuralima Boydstun resigned in July, and Wayne Worthen was hired in August to take over the project. Wayne began to develop an overall Project Manager Database used to manage the various databases required for the project.

Hatchery Operations Documentation

Clearwater Fish Hatchery

Brood Year 1993—Clearwater Fish Hatchery received 869,900 eyed eggs, Dworshak B-stock, from Dworshak National Fish Hatchery in April and May 1993 (George et al. 1995). Survival to the fry stage was 88.5% (769,862 fish). A total of 701,402 Dworshak B-stock steelhead were released: 50,027 fingerlings and 651,375 smolts (Appendix A. Table 1). Overall, eyed egg-to-release survival was 80.6%. Fingerlings, released as part of Idaho's Steelhead Supplementation Studies, were released into the South Fork Red River in September 1993. All fingerlings had a right ventral clip, and 5,000 were released with PIT tags. Dworshak B-stock smolts were released into Clear Creek (153,860), Crooked River (104,450), and the South Fork Clearwater River (393,065) in April and May 1994. There were 191,635 smolts released with CWTs and 1,364 fish released with PIT tags. Fish were tagged with CWTs in August 1993 and tagged with PIT tags in August 1993 and in March 1994. Fish were released in good health; no disease epizootic were reported.

A total of 136,476 green eggs, Selway B-stock, were collected during spawning operations at the Kamiah ponds (George et al. 1995). Of these, 93.2% (127,162) developed to the eyed stage. Survival to the fry stage was 86.8% (118,515). All Selway B-stock steelhead had the right pelvic fin

excised in September 1993. Adipose fins were not excised. A total of 71,566 Selway B-stock steelhead smolts were released into Crooked River in April and May 1994. Green egg-to-release survival was 52.4%. The relatively low egg-to-smolt survival was attributed to heavy bird predation during rearing. A total of 300 steelhead were tagged with PIT tags; no fish were tagged with CWTs. Selway B-stock smolts were PIT-tagged in April 1994.

Dworshak B- and Selway B-stock steelhead released at Crooked River were acclimated at the site for 10 days (April 18-28). Volitional release started on April 29, and all steelhead were forced out of the raceways on May 3, 1994.

Of the 1,364 Dworshak B-stock smolts released with PIT tags, 55.3% (754) were interrogated at downstream dams (Table 1). Interrogation rates, by PIT tag file, ranged from 33% to 86.4% (Table 1). The median travel time to Lower Granite Dam for PIT-tagged fish, by PIT tag file, ranged from 6.9 to 13.2 days (Table 1).

For an unknown reason, the steelhead released at two locations, Stites and the Red House Hole, had some fish that were PIT-tagged in August 1993 and some fish that were PIT-tagged in March 1994. That is, there were some fish released at each site that were tagged on different dates. Interestingly, for both release locations, the fish tagged in March (PIT tag files DAC94075.C*E) were interrogated at higher rates than the fish tagged in August (PIT tag files DAC93237.C*E) (Table 1). Moreover, for both release locations, the fish tagged in August had shorter median travel times than the fish tagged in March (Table 1). The lower interrogation rates for fish tagged in August could be due to undocumented tag loss or undetected PIT-tagged mortalities at the hatchery. However, migrational differences between the two groups may also be associated with fish size at the time of tagging. Fish tagged in August averaged 74 mm fork length, whereas the steelhead tagged in March averaged 165 mm fork length. Although this subject requires additional study, these findings suggest that steelhead smolts should be tagged in the spring prior to release rather than in the fall.

Comparing interrogation rates for the two stocks of steelhead released at Crooked River, 33% of the Dworshak B-stock fish were interrogated at downstream dams while only 5.3% of the Selway B-stock smolts were detected (Table 1). In addition, the median travel time of the Dworshak B-stock fish was considerably shorter—13.2 days as compared to 41.3 days. Besides the stock difference, the other major distinguishing feature between the two groups was smolt size-at-release. The Dworshak B-stock steelhead averaged 8.92 fish per pound (fpp) whereas the Selway B-stock fish averaged 24.3 fpp (Appendix A. Table 1).

PIT tag interrogation results for steelhead fingerlings used for supplementation are outside the scope of this report. Contact Alan Byrne (IDFG) for information concerning Idaho's Steelhead Supplementation Studies.

<u>Brood Year 1994</u>—Clearwater Fish Hatchery received 905,000 eyed eggs, Dworshak B-stock, from Dworshak National Fish Hatchery in April and May 1994 (McGehee et al. In Press). Eggs were collected at Dworshak National Fish Hatchery during the Number 10 through 13 egg-takes. Survival to the fry stage was 96.5% (873,511 fish). A total of 136,603 of the Dworshak B-stock steelhead were put on a two-year rearing program. Outbreaks of Columnaris and Aeromonas occurred from August through October due to warm water temperatures. Aeromonas was treated by feeding 4% TM-100 feed.

Clearwater Fish Hatchery also received 67,516 eyed eggs for the National Biological Survey (NBS) to conduct a study which examined stock performance and stock productivity impacts of

hatchery supplementation. Eggs were comprised of wild Selway B-stock fish, hatchery Dworshak B-stock fish, and a cross between the two stocks. These fish were also reared on a two-year rearing program. See Rankin and McGehee (In Press) for details on egg collection and hatchery rearing and Rubin et al. (1994) for the experimental design of the NBS study.

Hagerman National Fish Hatchery

<u>Brood Year 1993</u>—Hagerman National Fish Hatchery received a total of 2,019,973 eyed steelhead eggs in May and June 1993 comprised of two different stocks: Sawtooth A-stock (1,014,960 eggs) and Pahsimeroi A-stock (1,005,013 eggs) (Hagerman National Fish Hatchery 1993). Survival rates from the eyed egg stage to the fry stage for the Sawtooth A- and Pahsimeroi A-stocks were 96.4% and 95.6%, respectively. A total of 338,000 fingerlings were transferred to Niagara Springs Fish Hatchery in November 1993. Survival rates from eyed eggs to smolt release for Sawtooth A- and Pahsimeroi A-stocks were 94.1% (955,217 fish) and 84.6% (563,951 fish), respectively. No major health problems were reported.

Adipose fins were excised from all fish in October and November 1993. In November, approximately 246,000 steelhead were tagged with CWTs and marked by excising the left pelvic fin. Fish were tagged with CWTs to estimate adult contribution to the fishery and to test the effects of acclimating smolts at Sawtooth Fish Hatchery. In February 1994, 1,302 steelhead were tagged with PIT tags to evaluate emigration survival and timing to downstream dams.

Hagerman National Fish Hatchery released 1,519,168 steelhead smolts into the Salmon and Little Salmon rivers between April 6 and 29, 1994 (Appendix A. Table 2). The total fish release included 955,217 Sawtooth A-stock and 563,951 Pahsimeroi A-stock steelhead. Fish were released at four locations: Sawtooth Fish Hatchery weir (773,134), Little Salmon River (328,163), Lemhi River (235,788), and Bruno's Landing (Salmon River) (182,083). Of the steelhead released at Sawtooth Fish Hatchery weir, 712,362 were acclimated in raceways at Sawtooth Fish Hatchery between 8 and 17 days prior to being released. Some of these fish were tagged with CWTs and PIT tags as part of the Sawtooth Acclimation Experiment. Acclimated fish were released volitionally on April 25; all fish were forced from the raceways on April 29. Overall, there were 245,567 fish released from Hagerman National Fish Hatchery with CWTs and 1,300 fish released with PIT tags.

Excluding the Acclimation Experiment, interrogation rates for groups of PIT-tagged fish ranged from 48% to 64% (Table 1). Median travel times to Lower Granite Dam for groups of PIT-tagged fish ranged from 14.3 to 26.6 days (Table 1). See *Results, Experimentation – Objective 2, Hagerman National Fish Hatchery, Acclimation Experiment* for PIT tag interrogation rates of acclimated and non-acclimated steelhead.

<u>Brood Year 1994</u>—Hagerman National Fish Hatchery received a total of 1,673,647 eyed steelhead eggs comprised of three different stocks: Sawtooth A-stock (593,953 eggs), Pahsimeroi A-stock (362,118 eggs), and Oxbow A-stock (717,576 eggs) (Hagerman National Fish Hatchery 1994). Survival rates from eyed eggs to the fry stage for the Sawtooth A-, Pahsimeroi A-, and Oxbow A-stocks were 92.5% (549,312 fish), 98.1% (355,234 fish), and 94.7% (679,699 fish), respectively.

Magic Valley Fish Hatchery

<u>Brood Year 1993</u>—Magic Valley Fish Hatchery received a total of 2,767,613 eyed steelhead eggs comprised of three different stocks: Dworshak B-stock (1,507,033 eggs), Pahsimeroi A-stock (1,081,500 eggs), and East Fork B-stock (179,080 eggs) (Ainsworth et al. 1995). Survival rates from the eyed egg stage to the fry stage were as follows: Dworshak B-stock—97.1%, Pahsimeroi A-stock—99.3%, and East Fork B-stock—99.4%. In July 1993, 392,300 surplus fry, Dworshak B-stock, were shipped to Salmon Falls (227,600 fry) and Oakley (164,700 fry) reservoirs. Overall survival to release was 83.5% (2,311,550 fish—includes 392,300 fry).

In October and November 1992, 2,166,596 steelhead were marked with adipose fin clips. In November, 341,249 fish were tagged with CWTs and marked by excising the left pelvic fin. Coded-wire tags were used to determine fishery contribution. In February 1993, 1,701 steelhead (not tagged with CWTs) were tagged with PIT tags.

Magic Valley Fish Hatchery released a total of 1,919,250 brood year 1993 steelhead smolts at five different locations between April 11 and 28, 1994 (Appendix A. Table 3). The total release included 807,220 Dworshak B-stock, 951,990 Pahsimeroi A-stock, and 160,040 East Fork B-stock steelhead. A total of 337,775 fish, marked with a left pelvic fin clip, were released with CWTs. In addition, 1,701 steelhead were released with PIT tags (Appendix A. Table 3). Infectious pancreatic necrosis virus was isolated in all three stocks of fish; however, few fish died of the virus. Hatchery personnel reported that the fish were healthy at liberation.

Interrogation rates for groups of PIT-tagged fish ranged from 33% to 68% (Table 1). Median travel times to Lower Granite Dam for groups of PIT-tagged fish ranged from 11.5 to 24.1 days (Table 1). Overall, 49.4% of the PIT-tagged fish were interrogated at downstream dams.

<u>Brood Year 1994</u>—Magic Valley Fish Hatchery received a total of 2,396,340 eyed eggs comprised of three different stocks of steelhead: Dworshak B-stock (1,520,160 eggs), Pahsimeroi A-stock (800,785 eggs), and East Fork B-stock (75,395 eggs) (Moore et al. 1996). Overall survival to the fry stage was 80.5% (1,929,210 fish).

Migration Conditions

Snake River inflow (mean inflow) at Lower Granite Dam during the Peak and Extended periods in 1994 was 64.1 and 77.5 thousand cubic feet per second (kcfs), respectively (Table 2). With the exception of 1993, flows recorded during these periods were the largest since 1989.

Comparing Snake River flow conditions for the three brood years of steelhead that returned to Idaho as adults during this reporting period (broods 1989, 1990, and 1991), the 1989 brood, which emigrated in 1990, had the largest river discharge during the Peak migration period (Table 2). The 1990 brood, which emigrated in 1991, had the lowest river discharge during the Peak migration period. However, of the three broods, the 1990 brood had the largest river discharge during the Extended migration period (Table 2).

Migration Timing and Juvenile Survival

A total of 4,665 steelhead smolts were released with PIT tags in 1994: Clearwater Fish Hatchery—1,664, Hagerman National Fish Hatchery—1,300, and Magic Valley Fish Hatchery—1,701 (Table 1). Overall, 48.4% (2,256) of the PIT-tagged fish were interrogated at downstream dams. Interrogation rates of PIT-tagged steelhead, by PIT tag file, ranged from 5.3% to 73.7% (Table 1). Median travel times to Lower Granite Dam for PIT-tagged steelhead, by PIT tag file, ranged from 6.9 to 41.3 days (Table 1). Most of the steelhead tagged with PIT tags were interrogated at Lower Granite Dam between April 20 and May 15 (Figure 1). Flow conditions for the Snake River at Lower Granite Dam during this time period ranged from approximately 65 to 90 kcfs.

Adult Returns

The Harvest Monitoring Project (Ball 1997) estimated that Magic Valley Fish Hatchery and Hagerman National Fish Hatchery returned 7,534 steelhead to Idaho in 1993-1994 (Table 3). Ball (1997) estimated that 5,811 steelhead were harvested in Idaho's sport fishery and 1,723 steelhead returned to hatchery racks or escaped to spawn naturally. These estimates do not include tributary and mainstem strays or pre-spawning mortalities. The 1993-1994 steelhead return included 1-ocean, 2-ocean, and 3-ocean fish. The number of steelhead smolts released and the estimated number of adults that returned are compared to facility design production targets and projected adult return goals in Table 4. Hagerman National Fish Hatchery and Magic Valley Fish Hatchery achieved a minimum of 30% of their combined adult return goals (Table 4). Adult return estimates include only steelhead that returned to hatchery weirs, steelhead that were harvested in Idaho's sport fishery, and steelhead that escaped to spawn naturally.

Fisheries Contribution

Ball (1997) estimated that 5,811 LSRCP-reared hatchery steelhead were harvested during the 1993-1994 Idaho sport fishing season. See Ball (1997) for creel survey methods and results.

Hatchery Weirs

<u>Sawtooth Fish Hatchery Weir</u>—Hatchery fish returning to the Sawtooth weir in 1994 were released at the weir in 1991 and 1992 (brood years 1990 and 1991, respectively). Smolts were reared at Hagerman National and Magic Valley fish hatcheries prior to being trucked to Sawtooth Fish Hatchery and released.

A total of 338 adult steelhead (A-strain), comprised of 174 males (51.5%) and 164 females (48.5%), returned to the Sawtooth Fish Hatchery weir between March 16 and May 9, 1994 (Table 5) (Coonts 1994). The male component of the run was comprised of 173 hatchery-origin fish and one natural-origin (unmarked) fish; the female component was made up of 159 hatchery-origin fish and 5 natural-origin fish. All of the natural-origin fish and 54 of the hatchery-origin fish (32 males and 22 females) were released to spawn naturally. A total of 136 females and 141 males were spawned on seven different dates to yield 725,205 green eggs. A total of 660,989 eggs (91.1%) developed to the eyed stage.

<u>East Fork Salmon River Weir</u>—Hatchery fish returning to the East Fork Salmon River weir in 1994 were released at the weir in 1990, 1991, and 1992 (brood years 1989, 1990, and 1991,

respectively). Smolts released at the East Fork Salmon River weir were reared at Hagerman National and Magic Valley fish hatcheries.

A total of 73 adult steelhead (B-strain), comprised of 43 males (58.9%) and 30 females (41.1%), returned to the East Fork Salmon River weir between April 5 and May 4, 1994 (Table 6) (Coonts 1994). The male component of the run was comprised of 40 hatchery-origin fish (93%) and 3 natural-origin (unmarked) fish (7%); the female component was comprised of 25 hatchery-origin fish (83.3%) and 5 natural-origin fish (16.7%). All of the natural-origin fish and three male hatchery-origin fish were released to spawn naturally. Twenty-five females and 37 males were spawned on three different dates and produced 103,100 green eggs. A total of 76,087 eggs (73.8%) developed to the eyed stage.

Smolt-to-Adult Return Rates

Clearwater Fish Hatchery

Clearwater Fish Hatchery became operational in 1992. The first steelhead smolts were released in April 1993 (brood year 1992). Adults resulting from this release will return in 1995, 1996, and 1997.

Hagerman National Fish Hatchery

The 1993-1994 steelhead return included three age classes of fish which were released from Hagerman National Fish Hatchery in 1990 (brood year 1989), 1991 (brood year 1990), and 1992 (brood year 1991). Brood year 1989, 1990, and 1991 steelhead returned as 3-ocean, 2-ocean, and 1-ocean fish, respectively. The Harvest Monitoring Project estimated that 3,088 of the adult steelhead that returned to Idaho in 1993-1994 were reared at Hagerman National Fish Hatchery (Table 3). This equals 22.7% of the hatchery's adult return goal (13,600 fish). The adult return goal for Hagerman National Fish Hatchery was calculated based on the release of 2,400,000 smolts. However, smolt production for brood years 1989, 1990, and 1991 was 60% of the production goal (Table 4).

A total of 1,439,266 steelhead smolts were released from Hagerman National Fish Hatchery in 1990 (brood year 1989). An estimated 6,137 adult steelhead returned to Idaho from these smolts to yield a smolt-to-adult return rate (SAR) of 0.43% (Appendix B. Table 1). In 1991, 1,436,910 brood year 1990 steelhead smolts were released from Hagerman National Fish Hatchery (Appendix B. Table 2). The 2-ocean adult component for this brood returned in 1993-1994. To date, an estimated 5,269 adult steelhead have returned to Idaho from this brood to yield a SAR of 0.37%. In 1992, Hagerman National Fish Hatchery released 1,448,155 steelhead smolts from the 1991 brood (Appendix B. Table 3). Adult return data for this brood, specifically the 2- and 3-ocean components, are incomplete at this time.

The final adult steelhead from the 1989 (B-strain) and 1990 (A-strain) broods returned to Idaho during this reporting period. Smolt-to-adult return rates were calculated for coded-wire-tagged smolts, by stock and release site (Table 7). Dworshak B-stock steelhead smolts, brood year 1989, released into the Little Salmon River had a SAR of 1.28%. Large and normal size Sawtooth

A-stock steelhead smolts, brood year 1990, released at Sawtooth Fish Hatchery weir had smolt-to-adult return rates of 0.59% and 0.46%, respectively.

Magic Valley Fish Hatchery

The 1993-1994 steelhead return included three age classes of fish which were released from Magic Valley Fish Hatchery in 1990 (brood year 1989), 1991 (brood year 1990), and 1992 (brood year 1991). Brood year 1989, 1990, and 1991 steelhead returned as 3-ocean, 2-ocean, and 1-ocean fish, respectively. The Harvest Monitoring Project estimated that 4,446 of the adult steelhead that returned to Idaho in 1993-1994 were reared at Magic Valley Fish Hatchery (Table 3). This equals 38.1% of the hatchery's adult return goal (11,660 fish).

A total of 2,122,900 steelhead smolts were released in 1990 (brood year 1989) from Magic Valley Fish Hatchery (Appendix C. Table 1). An estimated 6,928 of these smolts returned to Idaho as adults (Appendix C. Table 1). The SAR for brood year 1989 was 0.33%. A total of 2,062,000 brood year 1990 steelhead smolts were released from Magic Valley Fish Hatchery in 1991 (Appendix C. Table 2). An estimated 7,256 of these smolts returned to Idaho as adults. The 3-ocean adult component of brood year 1990 is not complete. To date, the SAR for brood year 1990 is 0.35%. In 1992, Magic Valley Fish Hatchery released 2,160,400 brood year 1991 steelhead smolts (Appendix C. Table 3). Adult return data, specifically the 2- and 3-ocean components, for brood year 1991 are incomplete at this time.

The final adult steelhead from the 1989 (B-strain) and 1990 (A-strain) broods returned to Idaho during this reporting period. Smolt-to-adult return rates were calculated for coded-wire-tagged smolts, by stock and release site (Table 7). Smolt-to-adult return rates for Dworshak B- and East Fork B-stock steelhead smolts, brood year 1989, released into the East Fork Salmon River were 0.16% and 0.23%, respectively. Pahsimeroi A-stock steelhead smolts, brood year 1990, released into the Little Salmon River had a SAR of 0.35%.

Experimentation

Clearwater Fish Hatchery

<u>Cover Experiment</u>—Adults will return in 1995, 1996, and 1997. See Rhine et al. (In Press) for PIT tag results.

Hagerman National Fish Hatchery

<u>Size-at-Release Experiment</u>—The 2-ocean component of the 1990 brood returned during this reporting period. The total recoveries for the 1990 brood included 129 adults from large size smolts and 93 adults from regular size smolts (Table 8). Adult returns for the large size smolts were composed of 40% (51) females and 60% (78) males, whereas adults that returned from the regular size smolts were composed of 51% (47) females and 49% (46) males. Seventy-nine percent of the adults recovered from the large size smolts returned as 1-ocean fish, whereas 83% of the adults from the regular size smolts returned as 1-ocean fish.

The 1-ocean component of the 1991 brood returned during this reporting period. The total recoveries for the 1991 brood included ten adults from large size smolts and six adults from regular size smolts (Table 8). Complete results of this study will be reported under a separate title.

Acclimation Experiment—For brood year 1993, a total of 101 (33.9%) of the PIT-tagged acclimated steelhead were interrogated at downriver dams, as compared to 129 (42.7%) of the non-acclimated fish (Table 1). Interrogation rates were not significantly different (P > 0.05) among replicate raceways for each group. Therefore, data from replicate raceways, within group, were pooled and tested for differences using chi-square analysis. Significantly ($\chi^2 = 4.23$, P = 0.040) more of the non-acclimated steelhead smolts were interrogated as compared to the acclimated fish. Median travel times to Lower Granite Dam ranged from 11.3 to 17.8 days for the acclimated group and from 11.8 to 13.3 days for the non-acclimated group. Travel time to Lower Granite Dam was not significantly (P = 0.08) different between groups. Adult returns for brood year 1993 will be complete in 1997. Complete results of this study will be reported under a separate title.

The 1-ocean component of the 1991 brood returned during this reporting period. The total recoveries for the 1991 brood included eight adults from the acclimated smolts and six adults from non-acclimated smolts (Table 9). Adults from the acclimated group were all males, whereas 33.3% (2) of the adults from the non-acclimated group were males. Complete results of this study will be reported under a separate title.

LITERATURE CITED

- Ainsworth, B., D. May, D. Aplanalp, and K. Hills. 1995. Magic Valley Hatchery 1993 brood year report. Idaho Department of Fish and Game. Boise, Idaho.
- Ball, K. 1994. Evaluation of the hatchery-wild composition of Idaho salmon and steelhead harvest. United States Fish and Wildlife Service-Lower Snake River Fish and Wildlife Compensation Plan (October 1, 1991 to December 31, 1992). Idaho Department of Fish and Game. Boise, Idaho.
- Ball, K. 1996. Evaluation of the hatchery-wild composition of Idaho salmon and steelhead harvest. United States Fish and Wildlife Service-Lower Snake River Fish and Wildlife Compensation Plan (October 1, 1992 to December 31, 1993). Idaho Department of Fish and Game. Boise, Idaho.
- Ball, K. 1997. Evaluation of the hatchery-wild composition of Idaho salmon and steelhead harvest. United States Fish and Wildlife Service-Lower Snake River Fish and Wildlife Compensation Plan (October 1, 1993 to December 31, 1994). Idaho Department of Fish and Game. Boise, Idaho.
- Cannamela, D. A. 1992. Fish hatchery evaluations—Idaho. United States Fish and Wildlife Service-Lower Snake River Fish and Wildlife Compensation Plan Contract No. 14-16-0001-90502 (October 1, 1989 to September 30, 1990). Idaho Department of Fish and Game. Boise, Idaho.
- Columbia River Basin PIT Tag Information System. 1997. Pacific States Marine Fisheries Commission. Gladstone, Oregon.
- Coonts, P. J. 1994. Sawtooth Fish Hatchery and East Fork Satellite 1994 steelhead run report. Idaho Department of Fish and Game. Boise, Idaho.
- George, B., J. McGehee, and D. Munson. 1995. Clearwater Fish Hatchery 1992 chinook brood year and 1993 steelhead brood year reports. Idaho Department of Fish and Game. Boise, Idaho.
- Hagerman National Fish Hatchery. 1993. Hagerman National Fish Hatchery annual report, fiscal year 1993. United States Fish and Wildlife Service. Hagerman, Idaho.
- Hagerman National Fish Hatchery. 1994. Hagerman National Fish Hatchery annual report, fiscal year 1994. United States Fish and Wildlife Service. Hagerman, Idaho.
- McGehee, J., B. George, S. Patterson, and J. Rankin. In Press. Clearwater Fish Hatchery brood year 1994 chinook, brood year 1994 steelhead, and brood year 1995 steelhead reports. Idaho Department of Fish and Game. Boise, Idaho.
- Moore, B., D. May, K. Hills, and M. Olson. 1996. Magic Valley Hatchery 1994 brood year report. Idaho Department of Fish and Game. Boise, Idaho.

- Petrosky, C. E. 1991. Influence of smolt migration flows on recruitment and return rates of Idaho spring chinook. Idaho Department of Fish and Game. Boise, Idaho.
- Rankin, J. and J. McGehee. In Press. Clearwater Fish Hatchery Selway steelhead report 1994 and Crooked River steelhead report 1994. Idaho Department of Fish and Game. Boise, Idaho.
- Rhine, T. D., R. S. Osborne, and K. A. Stevens. In Press. Steelhead fish hatchery evaluations— Idaho. United States Fish and Wildlife Service-Lower Snake River Fish and Wildlife Compensation Plan Contract No. 14-48-0001-93500 (October 1, 1992 to September 30, 1993). Idaho Department of Fish and Game. Boise, Idaho.
- Rubin, S., L. Wetzel, and R. Reisenbichler. 1994. Performance/stock productivity impacts of hatchery supplementation. Progress Report 16 March to 15 June 1994. United States Department of Interior, National Biological Survey. Seattle, Washington.
- SYSTAT, Inc. 1992. SYSTAT for Windows: statistics, version 5 edition. Evanston, Illinois.

Table 1. Number of unique PIT tag interrogations of LSRCP steelhead smolts, by PIT tag file, at Lower Granite Dam (GRJ), Little Goose Dam (GOJ), Lower Monumental Dam (LMJ), and McNary Dam (MCJ) for the 1994 migration period. A total of 4,665 PIT-tagged steelhead were released from Hagerman National Fish Hatchery, Magic Valley Fish Hatchery, and Clearwater Fish Hatchery between April 5 and May 3, 1994. Median travel time is to Lower Granite Dam.

						N	umber /	Perce	nt Inte	rrogate	ed			Median
		Rel.	No.	G	RJ	G	OJ	LN	ЛJ	M	CJ	TO	TAL	Travel Time
File Name	Release Site	Date	Rel.	No.	%	No.	%	No.	%	No.	%	No.	%	(Days)
MAGIC VALLEY FIS	H HATCHERY													
Pahsimeroi A-stock														
Hazard Creek														
DAC94045.M14	Hazard Cr.	4/26	100	41	41.0	10	10.0	2	2.0	5	5.0	58	58.0	13.4
DAC94045.M16	Hazard Cr.	4/26	100	52	52.0	12	12.0	1	1.0	3	3.0	68	68.0	12.9
Hazard Creek Total			200									126	63.0	
Dworshak B-stock														
Hazard Creek														
DAC94044.M4E	Hazard Cr.	4/28	100	46	46.0	6	6.0	0	0.0	0	0.0	52	52.0	12.0
DAC94044.M4W	Hazard Cr.	4/28	100	39	39.0	9	9.0	1	1.0	3	3.0	52	52.0	12.1
DAC94044.M7E	Hazard Cr.	4/28	100	35	35.0	10	10.0	3	3.0	4	4.0	52	52.0	11.5
Hazard Creek Total			300									156	52.0	
Pahsimeroi A-stock														
Pahsimeroi Weir														
DAC94045.M9E	Pahsimeroi Weir	4/19	150	63	42.0	9	6.0	4	2.7	4	2.7	80	53.3	18.4
DAC94045.M11	Pahsimeroi Weir	4/19	151	74	49.0	13	8.6	3	2.0	3	2.0	93	61.6	16.3
Pahsimeroi Weir To	tal		301									173	57.5	
Dworshak B-stock														
East Fork Salmon Riv														
DAC94044.M1E	E.F. Salmon R.	4/15	100	30	30.0	6	6.0	3	3.0	0	0.0	39	39.0	21.6
DAC94044.M2E	E.F. Salmon R.	4/15	100	39	39.0	6	6.0	1	1.0	0	0.0	46	46.0	23.7
DAC94044.M3E	E.F. Salmon R.	4/15	100	36	36.0	3	3.0	4	4.0	1	1.0	44	44.0	17.9
East Fork Salmon R	iver Total		300									129	43.0	

Table 1. Continued.

Table 1. Continued.						N	lumber	/ Perce	nt Inte	rrogate	ed			Median
		Rel.	No.	G	RJ	G	OJ	LN	/IJ	М	CJ	TO	TAL	Travel Time
File Name	Release Site	Date	Rel.	No.	%	No.	%	No.	%	No.	%	No.	%	(Days)
East Fork B-stock														
East Fork Salmon R														
DAC94045.M7W	E.F. Salmon R.	4/16	100	41	41.0	8	8.0	1	1.0	0	0.0	50	50.0	20.5
DAC94045.M8E	E.F. Salmon R.	4/16	100	26	26.0	4	4.0	2	2.0	1	1.0	33	33.0	24.1
DAC94045.M8W	E.F. Salmon R.	4/16	100	41	41.0	8	8.0	3	3.0	3	3.0	55	55.0	21.2
East Fork Salmon I	River Total		300									138	46.0	
Dworshak B-stock														
Slate Creek (upper S	<u>Salmon River)</u>													
DAC94044.M5E	Slate Cr.	4/18	150	53	35.3	8	5.3	3	2.0	0	0.0	64	42.7	22.6
DAC94044.M6E	Slate Cr.	4/18	150	46	30.7	6	4.0	1	0.7	1	0.7	54	36.0	20.8
Slate Creek Total			300									118	39.3	
MAGIC VALLEY GR	RAND TOTAL		1,701									840	49.4	
	ONAL FISH HATCH	<u>ERY</u>												
Sawtooth A-stock														
	Freatment (Acclimate													
DAC94047.H54	Sawtooth FH	4/27	100	26	26.0	8	8.0	2	2.0	2	2.0	38	38.0	17.8
DAC94048.H55	Sawtooth FH	4/27	98	24	24.5	4	4.1	3	3.1	2	2.0	33	33.7	13.6
DAC94048.H56	Sawtooth FH	4/27	100	20	20.0	7	7.0	1	1.0	2	2.0	30	30.0	11.3
Treatment Group T	otal		298									101	33.9	
Sawtooth A-stock														
•	Control (Non-Acclima	,				4.0		_			4.0		40.4	44.0
DAC94047.H51	Sawtooth FH	4/29	102	30	29.4	10	9.8	3	2.9	1	1.0	44	43.1	11.8
DAC94047.H52	Sawtooth FH	4/29	100	30	30.0	9	9.0	6	6.0	2	2.0	47	47.0	12.6
DAC94047.H53	Sawtooth FH	4/29	100	25	25.0	3	3.0	5	5.0	5	5.0	38	38.0	13.3
Control Group Tota	al		302									129	42.7	
Pahsimeroi A-stock														
Hazard Creek	Hamand Or	4/05	400	40	40.0	_	0.0	-	- ^	4	4.0	00	00.0	440
DAC94048.H85	Hazard Cr.	4/25	100	43	43.0	8	8.0	5	5.0	4	4.0	60	60.0	14.3
DAC94048.H86	Hazard Cr.	4/25	100	42	42.0	11	11.0	7	7.0	2	2.0	62	62.0	14.5
Hazard Creek Total	I		200									122	61.0	

Table 1. Continued.

						N	lumber /	Perce	nt Inte	rrogate	ed			Median
		Rel.	No.	GI	٦J	G	OJ	LN	/IJ	M	CJ	TOT	ΓAL	Travel Time
File Name	Release Site	Date	Rel.	No.	%	No.	%	No.	%	No.	%	No.	%	(Days)
Sawtooth A-stock														
Salmon River (Brund	o's Landing)													
DAC94048.H74	Salmon R.	4/12	100	54	54.0	5	5.0	3	3.0	2	2.0	64	64.0	19.5
DAC94048.H75	Salmon R.	4/12	100	37	37.0	6	6.0	3	3.0	2	2.0	48	48.0	18.9
DAC94048.H76	Salmon R.	4/12	100	58	58.0	1	1.0	3	3.0	4	4.0	66	66.0	17.9
Salmon River Total			300									178	59.3	
Pahsimeroi A-stock														
<u>Lemhi River</u>														
DAC94048.H98	Lemhi R.	4/5	200	96	48.0	8	4.0	8	4.0	4	2.0	116	58.0	26.6
HAGERMAN NFH G	RAND TOTAL		1,300									646	49.7	
CLEARWATER FIS	H HATCHERY													
Dworshak B-stock														
Clear Creek														
DAC94075.C3W	Clear Cr.	5/3	300	188	62.7	18	6.0	10	3.3	5	1.7	221	73.7	6.9
Dworshak B-stock														
South Fork Clearwat	er River													
DAC94075.C4W	Cottonwood Cr.	4/25	200	115	57.5	14	7.0	7	3.5	2	1.0	138	69.0	8.5
DAC93237.C6E	Stites	4/26	158	72	45.6	9	5.7	2	1.3	1	0.6	84	53.2	8.8
DAC94075.C6E	Stites	4/26	39	18	46.2	7	17.9	1	2.6	1	1.0	27	69.2	12.4
Stites Total			197									111	56.3	
DAC93237.C7E	Red House Hole	4/26	156	78	50.3	8	5.2	5	3.2	1	0.6	92	59.0	7.7
DAC94075.C7E	Red House Hole	4/26	44	33	75.0	1	2.3	3	6.8	1	2.3	38	86.4	9.5
Red House Hole To	tal		200									130	65.3	
Dworshak B-stock														
Crooked River DAC93236.MIX	Crooked R.	5/1	467	76	16.3	52	11.1	17	3.6	9	1.9	154	33.0	13.2

			Number / Percent Interrogated											
		Rel.	No.	GI	₹J	G	OJ	LN	ΛJ	M	CJ	TO	ΓAL	Travel Time
File Name	Release Site	Date	Rel.	No.	%	(Days)								
Selway B-stock Crooked River														
TGC94102.SEL	Crooked R.	5/1	300	12	4.0	2	0.7	1	0.3	1	0.3	16	5.3	41.3
CLEARWATER GR	AND TOTAL		1,664									770	46.3	

Table 2. Snake River mean daily in-flow (thousand cubic feet per second) at Lower Granite Dam during the Peak and Extended chinook salmon smolt migration periods, 1977-1994. The migration periods are as defined by Petrosky (1991).

Year	Peak (04/15 - 05/05)	Extended (04/20 - 05/30)
1977	39.1	40.2
1978	85.4	95.8
1979	64.8	89.9
1980	87.5	102.9
1981	76.2	86.7
1982	116.8	131.6
1983	85.6	111.3
1984	121.9	146.1
1985	86.9	87.2
1986	93.4	105.7
1987	59.0	62.4
1988	55.1	64.2
1989	93.6	87.2
1990	63.8	66.4
1991	44.0	70.5
1992	54.2	57.3
1993	69.8	114.0
1994	64.1	77.5

Table 3. Estimated number of LSRCP steelhead that returned to Idaho in 1993-1994. The adult return in 1993-1994 included fish from three age classes. Steelhead were reared at Hagerman National and Magic Valley fish hatcheries. These estimates were prepared by the Idaho Department of Fish and Game's Harvest Monitoring Project and only include steelhead harvested in Idaho's sport fisheries, steelhead that returned to hatchery racks, and steelhead that escaped to spawn naturally. These are minimum estimates and do not include all tributary and mainstem strays or in-river pre-spawning mortalities.

HATCHERY	BROOD YEAR	3-OCEAN	2-OCEAN	1-OCEAN
Hagerman	1989	11		
Hagerman	1990		1,811	
Hagerman	1991			1,266
Estimated Return				3,088
Magic Valley	1989	47		
Magic Valley	1990		3,757	
Magic Valley	1991			642
Estimated Return				4,446
Total Return for B	oth Hatcheries			7,534

Table 4. Steelhead smolts released from Magic Valley and Hagerman National fish hatcheries that contributed to the 1993-1994 Idaho steelhead return. The number of steelhead smolts released and the estimated number of adults that returned were compared to facility design production targets and projected adult return goals.

	Releases Co	ntributing to	1993-1994 Adul	t Returns	_
Brood Year	Fish Hatchery	Number Released	Design Target	Percent of Target	1993-94 Adult Returns
1989	Magic Valley	2,122,900	2,000,000	106%	47
1989	Hagerman NFH	1,439,266	2,400,000	60%	11
	Total	3,562,166	4,400,000	81%	58
1990	Magic Valley	2,062,000	2,000,000	103%	3,757
1990	Hagerman NFH	1,436,910	2,400,000	60%	1,811
	Total	3,498,910	4,400,000	80%	5,568
1991	Magic Valley	2,160,400	2,000,000	108%	642
1991	Hagerman NFH	1,448,155	2,400,000	60%	1,266
	Total	3,608,555	4,400,000	82%	1,908
	Mean annual rele	ase as percen	t of target:	81%	
			Total adult retu	rn: ^a	7,534
			Adult return go	al:	25,260
			Percent of goal	l achieved:	30%

^a Estimate includes only steelhead harvested in Idaho's sport fisheries, steelhead that returned to hatchery racks, and off-site escapement. These are minimum estimates and do not include all tributary and mainstem strays or in-river pre-spawning mortalities.

Table 5. Summary of the 1994 A-strain steelhead trout return to the Sawtooth Fish Hatchery weir. The fish return included fish of hatchery and natural origin. Hatchery aging criteria, based on length, were used to determine agea. ND indicates "no data" (i.e., data were not available).

	HATCHERY ORIGIN n = 332											
		Ma	les n = 173		Females n = 159							
Age ^b	Trapped	Released	Spawned	Dead	Other	Trapped	Release	Spawne	Dead	Other		
							d	d				
1-ocean	85	ND	ND	0	0	46	ND	ND	0	ND		
2-ocean	88	ND	ND	0	0	113	ND	ND	0	ND		
Total	173	32°	141	0	0	159	22 ^c	136	0	1 ^d		

	NATURAL ORIGIN n = 6											
		M	ales n = 1	Females n = 5								
Age ^b	Trapped	Released	Spawned	Dead	Other	Trapped	Release	Spawne	Dead	Other		
							d	d				
1-ocean	1	1	0	0	0	0	0	0	0	0		
2-ocean	0	0	0	0	0	5	5	0	0	0		
Total	1	1 ^c	0	0	0	5	5°	0	0	0		

Total number trapped	338	Green egg number	725,205	
Trapping period	3/16 - 5/9/94	Eyed egg number	660,989	(91.1% eye - up)

^a Fish were aged using the following aging criteria:

<u>RUN</u>	SEX	LENGTH	AGE
Α	male	≤ 68 cm	1-ocean
Α	male	> 68 cm	2-ocean
Α	female	≤ 65 cm	1-ocean
Α	female	> 65 cm	2-ocean

b Hatchery fish classified as 1-ocean were released in 1992, brood year 1991. Hatchery fish classified as 2-ocean were released in 1991, brood

c Fish were released above the weir.
d The fish was killed, but the eggs were not used.

Summary of the 1994 B-strain steelhead trout return to the East Fork Salmon River weir. The fish return included fish of Table 6. hatchery and natural origin. Hatchery aging criteria, based on length, were used to determine agea. ND indicates "no data" (i.e., data were not available).

HATCHERY ORIGIN n = 65										
Males n = 40					Females n = 25					
Age⁵	Trapped	Release	Spawned	Dead	Other	Trapped	Release	Spawne	Dead	Othe
		d				· -	d	d		r
1-ocean	14	ND	ND	0	0	1	0	ND	0	0
2, 3-ocean	26	ND	ND	0	0	24	0	ND	0	0
Total	40	3 ^c	37	0	0	25	0	25	0	0

NATURAL ORIGIN n = 8										
Males n = 3					Females n = 5					
Age ^b	Trapped	Release	Spawned	Dead	Other	Trapped	Release	Spawne	Dead	Othe
		d					d	d		r
1-ocean	2	2	0	0	0	0	0	0	0	0
2, 3-ocean	1	1	0	0	0	5	5	0	0	0
Total	3	3 ^c	0	0	0	5	5 ^c	0	0	0

Total number trapped	73	Green egg number	103,100	
Trapping period	4/5 - 5/4/94	Eyed egg number	76,087	(73.8% eye - up)

^a Fish were aged using the following aging criteria:

<u>RUN</u>	<u>SEX</u>	<u>LENGTH</u>	<u>AGE</u>
В	male	≤ 73 cm	1-ocean
В	male	> 73 cm	2- or 3-ocean
В	female	≤ 68 cm	1-ocean
В	female	> 68 cm	2- or 3-ocean

b Ternale > 56 cm 2- or 3-ocean

b Hatchery fish classified as 1-ocean were released in 1992, brood year 1991. Hatchery fish classified as 2 or 3-ocean were released in 1991 and 1990, respectively, (brood years 1990 and 1989, respectively).

c Fish were released above the weir.

Table 7. Smolt-to-adult return rates of coded-wire-tagged steelhead smolts released from Hagerman National and Magic Valley fish hatcheries. The number of adults was estimated by Ball (1997) and only include steelhead harvested in Idaho's sport fisheries, steelhead that returned to hatchery racks, and steelhead that escaped to spawn naturally. These are minimum estimates and do not include all tributary and mainstem strays or in-river pre-spawning mortalities.

HATCHERY	STOCK	BROOD	RELEASE SITE	NUMBER CWT	NUMBER ADULTS	SAR (%)
Hagerman	DWOR B	1989	Little Salmon R.	42,322	542	1.28
Hagerman	SAW A	1990	Sawtooth FH	53,245	312	0.59
Hagerman	SAW A	1990	(large smolts) Sawtooth FH (regular smolts)	61,431	283	0.46
Magic	E.F. B	1989	E.F. Salmon R.	87,308	198	0.23
Magic	DWOR B	1989	E.F. Salmon R.	44,763	72	0.16
Magic	PAH A	1990	Little Salmon R.	65,997	229	0.35

Table 8. Total number^a of steelhead recovered with CWTs designating them as either large size^b or regular size^c. Recovered fish were released as smolts at the Sawtooth Fish Hatchery weir in 1991 (brood year 1990) and 1992 (brood year 1991). Recovery data were not available for the 2-ocean component of brood year 1991 (i.e., ND indicates "no data").

		1-C	cean	2-0			
Size Group	Brood Year	Number CWT	Male	Female	Male	Female	Total Return
Large	1990	53,245	66	36	12	15	129
Regular	1990	61,431	41	36	5	11	93
Large	1991	53,463	5	5	ND	ND	10
Regular	1991	45,646	4	2	ND	ND	6

^a Includes all Idaho fishery harvest returns, Idaho hatchery returns, and tributary stray recoveries

Table 9. Total number of steelhead recovered with CWTs designating them as either acclimated (ACC) or non-acclimated (NON-ACC). Recovered fish were released as smolts at the Sawtooth Fish Hatchery weir in 1992 (brood year 1991). ND indicates "No Data" (i.e., data were not available).

				n			
			1-C	cean	2-0	cean	
Exp. Group	Brood Year	Number CWT	Male	Female	Male	Female	Total Return
ACC	1991	55,632	8	0	ND	ND	8 ^b
NON-ACC	1991	45,646	4	2	ND	ND	6 ²
ACC	1992	65,865	ND	ND	ND	ND	ND
NON-ACC	1992	59,846	ND	ND	ND	ND	ND

^a Includes all Idaho fishery harvest returns, Idaho hatchery returns, and tributary stray recoveries

^b Large size steelhead averaged 241 mm (3.0 fish per pound) at time of release

^c Regular size steelhead averaged 221 mm (4.5 fish per pound) at time of release

^b Includes only 1-ocean returns




Figure 1. Number of unique interrogations of LSRCP juvenile steelhead at Lower Granite Dam, Washington, plotted with the average daily in-flow of the Snake River (at Lower Granite Dam) in 1994. A total of 4,665 PIT-tagged steelhead were released from Hagerman National, Magic Valley, and Clearwater fish hatcheries between April 5 and May 3, 1994. Thirty-seven percent (1,728) of the PIT-tagged fish were interrogated at Lower Granite Dam. Data for 16 fish fall outside of this date range and are not shown.

APPENDICES

Appendix A. Table 1. Release data and estimated adult returns for Clearwater Fish Hatchery summer steelhead trout, brood year 1993. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Dalassa	DW.	DW	011-		ntifying		0:	N	A	D-1 C	.	T-1-1	040
Release Site/Date	RW NO.	RW Total	Stock ID	Mark Type	CWT Code	Release Number	Size (FPP)	Marking Purpose	Age Ocean	Harvest	Composition Hatchery	Total Returns	SAR (%)
S.F. Clearwater R. 4/25/94	2W	28,417	DWOR B	CWT/LV/AD *LV/AD *AD	104731	20,424 741 7,252	8.3	Contribution	1 2 3	ND ND ND	ND ND ND	ND ND ND	ND ND ND
S.F. Clearwater R. 4/25-26/94	7E	52,419	DWOR B	CWT/LV/AD CWT/LV/AD/PIT *LV/AD *AD	104736	21,430 (200) 550 30,439	8.3	Contribution	1 2 3	ND ND ND	ND ND ND	ND ND ND	ND ND ND
S.F. Clearwater R. 4/25-26/94			DWOR B	**NON-CWT (Includes all *)		143,213	8.3						
S.F. Clearwater R. Milept. 18 above Stites 4/25-26/94	6E	52,103	DWOR B	CWT/LV/AD CWT/LV/AD/PIT *LV/AD *AD	104735	22,137 (197) 127 29,839	8.3	Contribution	1 2 3	ND ND ND	ND ND ND	ND ND ND	ND ND ND
S.F. Clearwater R. Milept. 18 above Stites 4/25-26/94			DWOR B	**NON-CWT (Includes all *)		82,165	8.3						
S.F. Clearwater R. Cottonwood Creek 4/25/94	4W	52,048	DWOR B	CWT/LV/AD CWT/LV/AD/PIT *LV/AD *AD AD/PIT	104734	21,086 (35) 630 30,332 (165)	9.99	Contribution	1 2 3	ND ND ND	ND ND ND	ND ND ND	ND ND ND
S.F. Clearwater R. Cottonwood Creek			DWOR B	**NON-CWT (Includes all *)		82,610							

4/25/94 Appendix A. Table 1. Continued.

Release Release Release Site Data Did Data Did Data Did Data Did Data Da						lentifying				_				
S.F. Clearwater R. Includes all release dates that have mark type indicated by ** DWOR B NON-CWT (Includes all "") 307,988 Production 1 ND ND ND ND ND ND ND									_	_		•		SAR
No No No No No No No No	Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
No		es that		DWOR B	(Includes all		307,988		Production					ND ND
Crooked River Pond 3E 52,188 DWOR B CWT.LV/AD 104732 21,779 8.92 Contribution 1 ND	have mark type indicate	d by **			,					3	ND	ND	ND	ND
A/29-5/3/94				TOTAL NO	N-CWT RELEA	- SE	307,988		TOTAL RETURN:		ND	ND	ND	ND
A/29-5/3/94	Crooked River Pond	3⊑	52 188	DWOR B	CWT/LV/AD	104732	21 770	8 02	Contribution	1	ND	ND	ND	ND
Crooked River Pond 4/29-5/3/94 Total cwt release total Non-cwt release		OL	02,100	DWORD		104702		0.02	Continuation	-				ND
A/29-5/3/94	4123-310134				LV/AD/PIT		(233)							ND
A/29-5/3/94	Crooked River Pond	7W	52,262	DWOR B	CWT/LV/AD	104737	20,883	8.92	Contribution	1	ND	ND	ND	ND
Crooked River Pond 12E SELWAY RV RV/PIT (300) Smolt Release 2 ND ND ND ND ND ND ND	4/29-5/3/94		,		*LV/AD					2	ND	ND	ND	ND
A/29-5/3/94 12W RV/PIT (300) Smolf Release 2 ND ND ND ND ND ND ND					LV/AD/PIT		(234)							ND
Crooked River Pond A/29-5/3/94 DWOR B NON-CWT 61,788 8.92 Production 1	Crooked River Pond			SELWAY				24.6					ND	ND
DWOR B	4/29-5/3/94	12W			RV/PIT		(300)		Smolt Release					ND
4/29-5/3/94 (Includes all *) (Includes all *)										3	ND	ND	ND	ND
TOTAL CWT RELEASE 42,662 TOTAL NON-CWT RELEASE 133,354 TOTAL SITE RELEASE 176,016 TOTAL RETURN: ND ND ND				DWOR B			61,788	8.92	Production					ND
TOTAL CWT RELEASE	4/29-5/3/94				(Includes all *)									ND
TOTAL NON-CWT RELEASE 133,354 176,016 TOTAL RETURN: ND ND ND ND										3	ND	ND	ND	ND
S.F. Red River V9 DWOR B RV 50,027 24.6 Supplementatio 1 ND ND ND ND						_								
S.F. Red River V9 DWOR B RV 50,027 24.6 Supplementatio 1 ND ND ND 9/1-2/93 RV/PIT (5,000) FingerFall Rel. 2 ND ND ND ND Alan Byrne's 3 ND ND ND ND TOTAL CWT RELEASE 0 TOTAL NON-CWT RELEASE 50,027						SE								
9/1-2/93 RV/PIT (5,000) FingerFall Rel. 2 ND ND ND Alan Byrne's 3 ND ND ND ND TOTAL CWT RELEASE 0 TOTAL NON-CWT RELEASE 50,027				TOTAL SIT	E RELEASE		176,016		TOTAL RETURN:		ND	ND	ND	ND
Alan Byrne's 3 ND ND ND TOTAL CWT RELEASE 0 TOTAL NON-CWT RELEASE 50,027		V9		DWOR B			·	24.6	n	•				ND
TOTAL CWT RELEASE 0 TOTAL NON-CWT RELEASE 50,027	9/1-2/93				RV/PIT		(5,000)							ND ND
TOTAL NON-CWT RELEASE 50,027				TOTAL CW	T RELEASE	_	0		,					
						SE	50,027							
TOTAL SITE RELEASE 50,027 TOTAL RETURN: ND ND ND							50,027		TOTAL RETURN:		ND	ND	ND	ND

Appendix A. Table 1. Continued

					entifying l	Marks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Return C	omposition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Clear Creek 5/3/94	1E	51,498	DWOR B	CWT/LV/AD *LV/AD *AD	104729	21,507 261 29,730	8.92	Contribution	1 2 3	ND ND ND	ND ND ND	ND ND ND	ND ND ND
Clear Creek 5/3/94	2E	51,411	DWOR B	CWT/LV/AD *LV/AD *AD	104730	21,562 240 29,609	8.92	Contribution	1 2 3	ND ND ND	ND ND ND	ND ND ND	ND ND ND
Clear Creek 5/3/94	3W	50,951	DWOR B	CWT/LV/AD CWT/LV/AD/PIT *LV/AD *AD AD/PIT	104733	20,827 (104) 360 29,764 (196)	8.92	Contribution	1 2 3	ND ND ND	ND ND ND	ND ND ND	ND ND ND
Clear Creek 5/3/94			DWOR B	NON-CWT (Includes all *)		89,964	8.92	Production	1 2 3	ND ND ND	ND ND ND	ND ND ND	ND ND ND
			TOTAL NO	VT RELEASE DN-CWT RELEAS TE RELEASE	SE	63,896 89,964 153,860		TOTAL RETURN:		ND	ND	ND	ND
TOTAL DWOR B-STO TOTAL DWOR B-STO TOTAL DWOR B-STO	CK NON	-CWT RE	_			191,635 509,767 701,402							
TOTAL SELWAY STO TOTAL SELWAY STO TOTAL SELWAY STO	CK NON	I-CWT RE				0 71,566 71,566							
TOTAL CWT RELEAS TOTAL NON-CWT RE TOTAL CLEARWATER	LEASE I	OR CLE		₹H		191,635 581,333 772,968							
TOTAL PIT TAGS						6,664							

Appendix A. Table 2. Release data and estimated adult returns for Hagerman National Fish Hatchery summer steelhead trout, brood year 1993. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Hager	man N	ational FH	Brood Y	ear: 1993									
				ld	entifying	Marks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Return C	Composition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Little Salmon R. 4/25/94	85	164,679	PAH A	CWT/LV/AD CWT/LV/AD/PI' *LV/AD *AD	104601 T	20,529 (100) 156 143,994	4.7	Contribution	1 2	ND ND	ND ND	ND ND	ND ND
Little Salmon R. 4/25/94	86	163,484	PAH A	CWT/LV/AD CWT/LV/AD/PI *LV/AD *AD	104602 T	20,365 (100) 275 142,844	4.5	Contribution	1 2	ND ND	ND ND	ND ND	ND ND
Little Salmon R. 4/25/94			PAH A	NON-CWT (Includes all *)		287,269		Production	1 2	ND ND	ND ND	ND ND	ND ND
			TOTAL N	WT RELEASE ION-CWT RELEA ITE RELEASE	ASE	40,894 287,269 328,163		TOTAL RETURN:		ND	ND	ND	ND
			IOIALO	TIL KLLLAGE		020,100		TOTAL KLIOKIN.		ND	ND	ND	ND
Lemhi River 4/6/94	98	235,788	PAH A	CWT/LV/AD CWT/LV/AD/PI *LV/AD *AD	104603 T	21,155 (200) 145 214,488	4.7	Contribution	1 2	ND ND	ND ND	ND ND	ND ND
Lemhi River 4/6/94			PAH A	NON-CWT (Includes all *)		214,633		Production	1 2	ND ND	ND ND	ND ND	ND ND
			TOTAL N	WT RELEASE ON-CWT RELEASE	ASE	21,155 214,633 235,788		TOTAL RETURN:	ND	ND	ND	ND	ND

Appendix A.	Table 2	Continued
ADDELIUIA A.	I abic 2.	COHUHUCU

тррепакт. тав				Id	entifying	Marks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		composition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Salmon River	51-3	61,321	SAW A	CWT/LV/AD	104628	60,772	4.7	Acclimation Control	1	ND	ND	ND	ND
Sawtooth Weir 4/29/94				CWT/LV/AD/PI *LV/AD	Γ	(302) 549			2	ND	ND	ND	ND
Salmon River Sawtooth Weir 4/15/94	54-6	711,813	SAW A	CWT/LV/AD CWT/LV/AD/PI *LV/AD *AD	104629 Г	60,929 (298) 507 650,377	4.4	Acclimation Test	1 2	ND ND	ND ND	ND ND	ND ND
Salmon River Sawtooth Weir 4/15-29/94			SAW A	NON-CWT (Includes all *)		651,433		Production	1 2	ND ND	ND ND	ND ND	ND ND
			TOTAL C	WT RELEASE		121,701							
				ON-CWT RELEA	SE	651,433							
			TOTAL S	ITE RELEASE		773,134		TOTAL RETURN:		ND	ND	ND	ND
Bruno Landing 4/12/94	74-6	182,083	SAW A	CWT/LV/AD CWT/LV/AD/PI *LV/AD *AD	104626 Г	61,817 (300) 778 119,488	4.7	Contribution	1 2	ND ND	ND ND	ND ND	ND ND
Bruno Landing 4/12/94			SAW A	NON-CWT (Includes all *)		120,266		Production	1 2	ND ND	ND ND	ND ND	ND ND
			TOTAL N	WT RELEASE ON-CWT RELEA ITE RELEASE	SE	61,817 120,266 182,083		TOTAL RETURN:		ND	ND	ND	ND
TOTAL PAH A-S TOTAL PAH A-S TOTAL PAH A-S	TOCK	NON-CWT	RELEASE	Ē		62,049 501,902 563,951							
TOTAL SAW A-S TOTAL SAW A-S TOTAL SAW A-S	тоск	NON-CW	T RELEASE	Ē		183,518 771,699 955,217							
TOTAL CWT REI		-		-	н	245,567 1,273,601							

TOTAL HAGERMAN NATIONAL FH RELEASE
TOTAL PIT TAGS

1,519,168 1,300

Appendix A. Table 3. Release data and estimated adult returns for Magic Valley Fish Hatchery summer steelhead trout, brood year 1993. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Magic Valle	ey FH	Brood '	Year: 1993										
, ,	-			le	dentifying	Marks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Return C	Composition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
E. Fk. Salmon River	1E	63,970	DWOR B	CWT/LV/AD	104721	20,443	5.6	Contribution	1	ND	ND	ND	ND
4/13-14/94				*LV/AD		181			2	ND	ND	ND	ND
				*AD		43,346			3	ND	ND	ND	ND
				AD/PIT		(100)							
E. Fk. Salmon River	2E	59,925	DWOR B	CWT/LV/AD	104722	20,974	5.5	Contribution	1	ND	ND	ND	ND
4/14/94				*LV/AD		319			2	ND	ND	ND	ND
				*AD		42,737			3	ND	ND	ND	ND
				AD/PIT		(100)							
E. Fk. Salmon River	3E	64,400	DWOR B	CWT/LV/AD	104710	21,296	4.4	Contribution	1	ND	ND	ND	ND
4/15/94		,		*AD		43,104			2	ND	ND	ND	ND
				AD/PIT		(100)			3	ND	ND	ND	ND
E. Fk. Salmon River 4/11-12/94	1-3W	164,740	DWOR B	*AD		164,740							
E. Fk. Salmon River	7W	54,720	E. FK. B	CWT/LV/AD	104711	20,648	4.8	Contribution	1	ND	ND	ND	ND
4/12/94		,		*LV/AD		179			2	ND	ND	ND	ND
				*AD		33,893			3	ND	ND	ND	ND
				AD/PIT		(100)							
E. Fk. Salmon River	8W	54,400	E. FK. B	CWT/LV/AD	104712	21,377	4.8	Contribution	1	ND	ND	ND	ND
4/12-13/94				*LV/AD		201			2	ND	ND	ND	ND
				*AD		32,822			3	ND	ND	ND	ND
				AD/PIT		(100)							
E. Fk. Salmon River	8E	50,920	E. FK. B	CWT/LV/AD	104713	21,369	4.8	Contribution	1	ND	ND	ND	ND

4/15-16/94 Appendix A. Table 3.	Continuo	,d		*AD AD/PIT		29,551 (100)			2 3	ND ND	ND ND	ND ND	ND ND
Appendix A. Table 5.	Continue	u.		le	dentifying	Marks							
Release Site/Date	RW NO.	RW Total	Stock ID	Mark Type	CWT Code	Release Number	Size (FPP	Marking Purpose	Age Ocean	Return C Harvest	Composition Hatchery	Total Returns	SAR (%)
E. Fk. Salmon River 4/11-16/94			E. FK. B DWOR B	NON-CWT (Includes all *)		391,073		Production	1 2	ND ND	ND ND	ND ND	ND ND
				•					3	ND	ND	ND	ND
			TOTAL NO	WT RELEASE ON-CWT RELE TE RELEASE	- ASE	126,107 391,073 517,180		TOTAL RETU	RN:	ND	ND	ND	ND
L. Salmon River at Warm Spring 4/26/94	14E	58,000	PAH A	CWT/LV/AD *LV/AD *AD AD/PIT	104723	20,958 180 36,862 (100)	4.4	Contribution	1 2	ND ND	ND ND	ND ND	ND ND
L. Salmon River at Warm Spring 4/27/94	16E	56,730	РАН А	CWT/LV/AD *LV/AD *AD AD/PIT	104724	21,530 200 35,000 (100)	4.1	Contribution	1 2	ND ND	ND ND	ND ND	ND ND
L. Salmon River at Warm Spring 4/23-27/94	13-15E 13-16V		PAH A	*AD		352,820							
L. Salmon River at Warm Spring			PAH A	NON-CWT (Includes all *)		425,062		Production	1 2	ND ND	ND ND	ND ND	ND ND
4/23-27/94)									
			TOTAL NO	WT RELEASE ON-CWT RELE TE RELEASE	- ASE	42,488 425,062 467,550		TOTAL RETU	RN:	ND	ND	ND	ND
Slate Creek U. Salmon R. 4/14-18/94	5E 6E	120,215	DWOR B	CWT/LV/AD *LV/AD *AD AD/PIT	104925	62,510 625 57,080 (300)		Contribution	1 2 3	ND ND ND	ND ND ND	ND ND ND	ND ND ND

Slate Creek 5W U. Salmon R. 6W 4/12-20/94

DWOR B *AD

91,140

Appendix A. Table 3. Continued.

					dentifying								
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Return C	omposition	Total	SAF
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Slate Creek			DWOR B	NON-CWT		148,845		Production	1	ND	ND	ND	ND
U. Salmon R.				(Includes all *)					2	ND	ND	ND	ND
4/12-20/94)					3	ND	ND	ND	ND
			TOTAL C	WT RELEASE	_	62,510							
				ON-CWT RELE	ASE	148,845							
			TOTAL SI	TE RELEASE		211,355		TOTAL RETU	RN:	ND	ND	ND	ND
Pahsimeroi River	9E	60,580	PAH A	CWT/LV/AD	104725	21,294	4.7	Contribution	1	ND	ND	ND	ND
4/16-18/94				*LV/AD		113			2	ND	ND	ND	ND
				*AD AD/PIT		39,173 (150)							
Pahsimeroi River	11E	60,920	PAH A	CWT/LV/AD	104726	21,660	5.6	Contribution	1	ND	ND	ND	ND
4/19/94		,		*LV/AD		330			2	ND	ND	ND	ND
				*AD AD/PIT		38,930 (151)							
						, ,							
Pahsimeroi River 4/18-22/94	9-12W 10-12E	<u> </u>	PAH A	*AD		362,940							
Pahsimeroi River			PAH A	NON-CWT		441,486		Production	1	ND	ND	ND	ND
4/16-22/94				(Includes all *)					2	ND	ND	ND	ND
			TOTAL C	WT RELEASE	_	42,954							
				ON-CWT RELE	ASE	441,486				ND	ND	ND	ND
			IOTAL SI	TE RELEASE		484,440		TOTAL RETU	KN:	ND	ND	ND	ND
Hazard Creek	4W	64,770	DWOR B	CWT/LV/AD	104714	20,891	5.1	Contribution	1	ND	ND	ND	ND
Little Salmon River 4/22/94				*LV/AD *AD		297 43,582			2 3	ND ND	ND ND	ND ND	ND ND
4122134				AD		40,002			3	טוו	שוו	ND	טאו

				AD/PIT		(100)							
Hazard Creek Little Salmon River 4/25-28/94	4E	63,840	DWOR B	CWT/LV/AD *AD AD/PIT	104715	21,243 42,597 (100)	4.2	Contribution	1 2 3	ND ND ND	ND ND ND	ND ND ND	ND ND ND
Appendix A. Table 3. (Continue	ed.											
Release	RW	RW	Stock	Mark	lentifying CWT	Marks Release	Size	Marking	۸۵۵	Doturn C	Composition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP	Purpose	Age Ocean	Harvest	Hatchery	Returns	(%)
Oncorputo				1,700		- Trainison)	- Lapose	- Cocum	Tidi Voot	- Hatoriory	Ttotarrio	(70)
Hazard Creek Little Salmon River 4/28/94	7E	60,390	DWOR B	CWT/LV/AD *LV/AD *AD AD/PIT	104716	21,582 154 38,654 (100)	4.5	Contribution	1 2 3	ND ND ND	ND ND ND	ND ND ND	ND ND ND
Hazard Creek Little Salmon River 4/22-23/94	3W 6W	49,725	DWOR B	*AD		49,725	5.1						
Hazard Creek Little Salmon River			DWOR B	NON-CWT (Includes all *)		175,009		Production	1 2	ND ND	ND ND	ND ND	ND ND
4/22-28/94				,					3	ND	ND	ND	ND
			TOTAL NO	VT RELEASE DN-CWT RELEASE TE RELEASE	ASE	63,716 175,009 238,725		TOTAL RETU	RN:	ND	ND	ND	ND
Oakley Reservoir 7/14/94	1W	164,700	DWOR B	No Mark		164,700	183	Fry Outplants					
			TOTAL NO	VT RELEASE ON-CWT RELEASE TE RELEASE	ASE	0 164,700 164,700		TOTAL RETU	RN:	ND	ND	ND	ND
Salmon Falls Reservoir 7/13/94	V's	227,600	DWOR B	No Mark		227,600	228	Fry Outplants					

TOTAL CWT RELEASE

0

ယ
m
\sim

	TOTAL NON-CWT RELEASE	227,600					
	TOTAL SITE RELEASE	227,600	TOTAL RETURN:	ND	ND	ND	ND
TOTAL E. FK. B-STOCK CWT RELEA	ASE	63,394					
TOTAL E. FK. B-STOCK NON-CWT F	RELEASE	96,646					
TOTAL E. FK. B-STOCK RELEASE		160,040					
TOTAL DWOR B-STOCK CWT RELE	ASE	188,939					
TOTAL DWOR B-STOCK NON-CWT	RELEASE	1,010,581	(includes 392,300 fry re	eleased at Sa	ılmon Falls a	and Oakley	Res.)
TOTAL DWOR B- STOCK RELEASE		1,199,520	Smolt Release = 807,2	20			
Appendix A. Table 3. Continued.							
TOTAL PAH A-STOCK CWT RELEAS	SE	85,442					
TOTAL PAH A-STOCK NON-CWT RE	ELEASE	866,548					
TOTAL PAH A-STOCK RELEASE		951,990					
TOTAL CWT RELEASE FOR MAGIC	VALLEY FH	337,775					
TOTAL NON-CWT RELEASE FOR M	AGIC VALLEY FH	1,973,775					
TOTAL MAGIC VALLEY FH RELEAS	E	2,311,550					
TOTAL PIT TAGS		1,701					
TOTAL SMOLT RELEASE		1,919,250					

Appendix B. Table 1. Release data and estimated adult returns for Hagerman National Fish Hatchery summer steelhead trout, brood year 1989. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Hagerman NFH	Broo	d Year: 19	989										
-				ld	dentifying	Marks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Return C	omposition	Total	SAR
Site/Date	NO.	Total	ID	Type	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Sawtooth Weir	70	99,878	SAW A	CWT/LV/AD	104214	14,597	4.7	Contribution	1	76	19	113	0.77
4/5/90	71			*LV/AD *AD		147 85,134			2	12	6		
Sawtooth Weir	70	100,772	SAW A	CWT/LV/AD	104215	15,482	4.7	Contribution	1	48	12	98	0.63
4/5/90	71			*LV/AD *AD		156 85,134			2	33	5		
Sawtooth Weir	70	100,506	SAW A	CWT/LV/AD	104216	15,218	4.7	Contribution	1	60	15	78	0.51
4/5/90	71			*LV/AD		154			2	3	0		
				*AD		85,134							
Sawtooth Weir			SAW A	NON-CWT		255,859	4.7	Production	1	2,072	518	2,948	1.15
Includes all release dates that have Mark Type indicated by				(Includes all *)			(Includes all *)	2	314	44		
		•		WT RELEASE ON-CWT RELE	ACE	45,297 255,859							
				TE RELEASE	ASE	301,156		TOTAL RETU	RN:	2,618	619	3,237	1.07
Salmon R. at Shoup Bridge 4/12/90	92 93	67,001	SAW A	CWT/LV/AD *AD	104227	15,528 51,473	5	Contribution	1 2	24 0	17 0	41	0.26
4/12/90	95			AD		31,473			2	U	U		
Salmon R. at Shoup Bridge	92	66,669	SAW A	CWT/LV/AD	104228	15,196	5	Contribution	1	30	23	61	0.40
4/12/90	93			*AD		51,473			2	8	0		
Salmon R. at Shoup Bridge	92	66,576	SAW A	CWT/LV/AD	104229	15,104	5	Contribution	1	19	14	42	0.28
4/12/90	93			*AD		51,472			2	9	0		

Salmon R. at Shoup Bridge	92	SAW A NON-CWT	154,418	5	Production	1	179	119	355	0.23
Includes all release dates	93	(Includes all *)			(Includes all	2	57	0		
that					*)					
have Mark Type indicated by										

Appendix B. Table 1. Continued.

Appendix B. Table 1. Continu				lo	dentifying	Marks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		omposition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
			TOTAL NO	VT RELEASE ON-CWT RELE TE RELEASE	ASE	45,828 154,418 200,246		TOTAL RETU	RN:	326	173	499	0.25
Little Salmon R. at Hazard Cr. 4/30/90	48 49	131,854	DWOR B	CWT/LV/AD *LV/AD *AD	104230	14,841 49 116,964	4.4	Contribution	1 2 3	0 258 0	0 258 0	516	3.48
Little Salmon R. at Hazard Cr. 4/30/90	48 49	131,075	DWOR B	CWT/LV/AD *LV/AD *AD	104231	14,065 47 116,963	4.4	Contribution	1 2 3	0 4 0	0 4 0	8	0.06
Little Salmon R. at Hazard Cr. 4/30/90	49	130,423	DWOR B	CWT/LV/AD *LV/AD *AD	104232	13,416 44 116,963	4.4	Contribution	1 2 3	0 9 0	0 9 0	18	0.13
Little Salmon R. at Hazard Cr. 4/23/90	ND	ND	SAW A	*AD		80,465	4.2	Production	1 2	33 30	33 30	126	0.16
Little Salmon R. at Hazard Cr.	ND		DWOR B	NON-CWT		351,030	4.4	Production	1	0	0	332	0.09
Includes all release dates that				(Includes all *)			(Includes all	2	166	166		
have Mark Type indicated by								,	3	0	0		
*			TOTAL NO	VT RELEASE DN-CWT RELE TE RELEASE	ASE	42,322 431,495 473,817		TOTAL RETU	RN:	500	500	1,000	0.21
Salmon R. at Ellis Bridge 4/9-11/90	ND	ND		*AD VT RELEASE DN-CWT RELE	ASE	200,295 0 200,295	4.4	Production	1 2	252 256	168 24	700	0.35

			TOTAL SIT	E RELEASE		200,295		TOTAL RETU	IRN:	508	192	700	
Salmon R. at N. Fk Ramp 4/18-20/90	ND	ND	SAW A	*AD		199,602	4.3	Production	1 2	251 250	167 9	677	0.34
17 10 20,00			TOTAL CW	T RELEASE	<u>-</u> !	0			-	200	· ·		
			TOTAL NO			199,602							
			TOTAL SIT	E RELEASE		199,602		TOTAL RETU	IRN:	501	176	677	0.34
Appendix B. Table 1. Contir	nued.												
			_		Identifying	Marks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Return C	Composition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
East Fork Salmon R. 4/11/90	ND	ND	DWOR B	*AD		64,150	3.8	Production	1 2	3 9	0	24	
			TOTAL CW	T RELEASE	<u> </u>	0			3	11	0		
			TOTAL NO	N-CWT REL	EASE	64,150			-				
			TOTAL SIT	E RELEASE	!	64,150		TOTAL RETU	IRN:	23	1	24	0.04
T0T41 0414 4 0T0 01/ 041						04.405							
TOTAL SAW A-STOCK CW TOTAL SAW A-STOCK NOI			=			91,125 890,639							
TOTAL SAW A-STOCK NO	-	KELEAS	<u>=</u>			981,764	_	RETURN IS C	OMPLET	·F			
TOTAL DAW A GTOOK KEE	LACL					301,704		KETOKITIO	JOINI LLI	-			
TOTAL DWOR B-STOCK C	WT REL	EASE				42,322		RETURN					
TOTAL DWOR B-STOCK N	-		SE			415,180	_	GRAND TOTA	AL:	4,476	1,661	6,137	0.43
TOTAL DWOR B-STOCK R	ELEASE	•				457,502							
TOTAL CWT RELEASE FOI	R HAGE	RMAN N	FH			133,447							
TOTAL NON-CWT RELEAS	E FOR I	HAGERN	IAN NFH			1,305,819							
TOTAL HAGERMAN NFH R	ELEASE	Ē				1,439,266							

Appendix B. Table 2. Release data and estimated adult returns for Hagerman National Fish Hatchery summer steelhead trout, brood year 1990. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Hagerr	nan NF	H Brood	Year: 1990										
					dentifying								
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		Composition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Sawtooth Weir 4/13/91	71- 74	22,652	SAW A	CWT/LV/AD *LV/AD *AD	104333	21,050 213 1,389	2.9	Size at release Large size group	1 2	45 60	24 3	132	0.63
Sawtooth Weir 4/13/91	71- 74	21,580	SAW A	CWT/LV/AD *LV/AD *AD	104334	20,129 122 1,329	2.9	Size at release Large size group	1 2	55 26	27 5	113	0.56
Sawtooth Weir 4/15/91	74- 75	12,864	SAW A	CWT/LV/AD *AD	104335	12,066 798	2.6	Size at release Large size group	1 2	32 19	12 4	67	0.56
Sawtooth Weir 4/13/91	76- 78	326,644	SAW A	CWT/LV/AD *LV/AD *AD	104336	21,775 131 304,738	4.5	Size at release Regular size group	1 2	55 12	13 3	83	0.38
Sawtooth Weir 4/13/91	76- 78	305,400	SAW A	CWT/LV/AD *LV/AD *AD	104337	20,318 143 284,939	4.5	Size at release Regular size group	1 2	49 31	15 1	96	0.47
Sawtooth Weir 4/16/91	78- 80	290,660	SAW A	CWT/LV/AD *LV/AD *AD	104338	19,338 156 271,166	4.4	Size at release Regular size group	1 2	52 29	19 4	104	0.54
Sawtooth Weir 4/13-16/91	71- 80	985	SAW A	*AD/PIT		985 (489) (496)	ND	Size at release (Lg) Size at release (Reg)	1 2	2	1 0	4	0.41
Sawtooth Weir Includes all release	e dates	865,124 that	SAW A	NON-CWT (Includes all *)	865,124	2.9	Production (Includes all *)	1 2	2,173 1,013	682 132	4,000	0.46

have Mark Type indicated by * Appendix B. Table 2. Continued.

					lentifying								
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		Composition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Sawtooth Weir 10/5-17/90	ND	ND	SAW A	*AD	_	304,907	39.6	Fall release	1 2	0 0	0 0	0	0.00
				NT RELEASE		114,676							
			TOTAL N	ON-CWT RELE	ASE	1,170,031							
			TOTAL SI	TE RELEASE		1,284,707		TOTAL RETURN:		3,654	945	4,599	0.36
Little Salmon R.	19-	154,937	DWOR B	CWT/LV/AD	104332	19,831	4.4	Contribution	1	6	6	42	0.21
at Hazard Cr.	21			*LV/AD		384			2	15	15		
4/22/91				*AD		134,722			3	ND	ND		
Little Salmon R.	51	154,379	DWOR B	CWT/LV/AD	104339	19,813	4.4	Contribution	1	4	4	18	0.09
at Hazard Cr.				*LV/AD		241			2	5	5		
4/19/91				*AD		134,325			3	ND	ND		
Little Salmon R.	38	147,794	DWOR B	CWT/LV/AD	104340	18,877	4.5	Contribution	1	3	3	26	0.14
at Hazard Cr.				*LV/AD		554			2	10	10		
4/17/91				*AD		128,363			3	ND	ND		
Little Salmon R.	44	424	DWOR B	AD/PIT		424	4.5	Contribution	1	0	0	0	ND
at Hazard Cr.	45								2	0	0		
4/24/91									3	ND	ND		
Little Salmon R.	ND		DWOR B	AD		120,323	22.5	Excess Fingerling	1	27	27	54	0.04
at Hazard Cr.								Plant	2	0	0		
									3	ND	ND		
Little Salmon R. at Hazard Cr.			DWOR B	NON-CWT		398,589	4-4.5	Production	1	88	88	584	0.15
Includes all release have Mark Type in				(Includes all *))			(Includes all *)	2	204 ND	204 ND		
		~,	TOTAL C	NT RELEASE		58,521			-				
				ON-CWT RELE	ASE	518,912							
			TOTAL SI	TE RELEASE		577,433		TOTAL RETURN:		362	362	724	0.13
E. Fk Salmon R.	ND		UNK. B	AD		540,733	32.8	Excess Fingerling	1	29	4	33	0.01
(above trap)								Plant	2	0	0		
9/5-7/90									3	ND	ND		

Appendix B. Table 2. Continued.

				I	dentifying	Marks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Return C	Composition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
			TOTAL CW	/T RELEASE	_	0							
			TOTAL NO	N-CWT REL	EASE	540,733							
			TOTAL SIT	E RELEASE		540,733		TOTAL RETURN:		29	4	33	0.01
TOTAL SAW A-S	этоск с	WT RELE	EASE			114,676							
TOTAL SAW A-S	STOCK N	ON-CWT	RELEASE			1,170,031							
TOTAL SAW A-S	STOCK R	ELEASE				1,284,707	•						
								RETURN IS INCOM	PLETE				
TOTAL DWOR E						58,521							
TOTAL DWOR E						518,912	<u>-</u>	RETURN					
TOTAL DWOR E	3-STOCK	RELEAS	E			577,433		GRAND TOTAL:		4,045	1,311	5,356	0.22
TOTAL UNK B-S	TOCK C	WT DELE	ACE			0		(includes fingerling					
TOTAL UNK B-S						0 540,733		and fry releases)					
TOTAL UNK B-S			RELEASE			540,733	-						
TOTAL UNK B-S	JIOCK KI	LLLASE				340,733							
TOTAL CWT RE	LEASE F	OR HAGI	ERMAN NFH			173,197							
TOTAL NON-CW	T RELEA	SE FOR	HAGERMAN	NFH		2,229,676							
TOTAL HAGER	MAN NFH	RELEAS	Ε			2,402,873	•						
TOTAL PIT TAG	S					1,409							
TOTAL SMOLT	RELEASE	Ē.				1,436,910		RETURN (smolt releases	only)	3,989	1,280	5,269	0.37

Appendix B. Table 3. Release data and estimated adult returns for Hagerman National Fish Hatchery summer steelhead trout, brood year 1991. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Hagerman N	FH Bro	ood Year:	1991										
				le	dentifying	Marks		_					
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Return C	omposition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
E. Fk. Salmon R. at trap	ND	ND	DWOR B	*AD		302,335	ND	Production	1	292	14	306	0.10
4/6-9/92					_				2	ND	ND		
				WT RELEASE		0			3	ND	ND		
				ON-CWT RELE	EASE	302,335							
			TOTAL SI	TE RELEASE		302,335		TOTAL RETURN:		292	14	306	0.10
Little Salmon R.	ND	ND	DWOR B	CWT/LV/AD	104407	18,386	ND	Contribution	1	0	0	0	0.00
at Warm Springs Bridge				*LV/AD		316			2	ND	ND		
4/16,20/92				*AD		81,230			3	ND	ND		
Little Salmon R.	ND	ND	DWOR B	CWT/LV/AD	104408	19,450	ND	Contribution	1	0	0	0	0.00
at Warm Springs Bridge				*LV/AD		546			2	ND	ND		
4/14/92				*AD		85,930			3	ND	ND		
Little Salmon R.	ND	ND	DWOR B	CWT/LV/AD	104409	17,390	ND	Contribution	1	0	0	0	0.00
at Warm Springs Bridge				*LV/AD		457			2	ND	ND		
4/16/92				*AD		76,829			3	ND	ND		
Little Salmon R.	ND	ND	DWOR B	NON-CWT		245,308	ND	Production	1	0	0	0	0.00
at Warm Springs Bridge				(Includes all *)				2	ND	ND		
4/14,16,20,22/92					-				3	ND	ND		
				WT RELEASE	- 4 6 5	55,226							
				ON-CWT RELE	EASE	245,308		TOTAL RETURN:		0	0	0	0.00
			IOTAL SI	TE RELEASE		300,534		IOIAL REIORN:		U	U	U	0.00
Pahsimeroi ponds	61-70	223,406	PAH A	*AD		223,406	ND	Production	1	270	125	395	0.18
3/25-27/92					-				2	ND	ND		
				WT RELEASE		0							
			TOTAL N	ON-CWT RELE	EASE	223,406							

	Appendix B. Table 3.	Continued	1	TOTAL S	ITE RELEASE		223,406		TOTAL RETURN:		270	125	395	0.18
	Appendix B. Table 3.	Continued			1	dentifying	Marks							
_	Release Site/Date	RW NO.	RW Total	Stock ID	Mark Type	CWT Code	Release Number	Size (FPP)	Marking Purpose	Age Ocean	Return C Harvest	omposition Hatchery	Total Returns	SAR (%)
	Sawtooth Weir 3/13-21/92	71,84		РАН А	CWT/LV/AD *LV/AD *AD	104421	17,955 152 149,183	4.8	Acclimation Experimental Group Trucked to SAWT on 3/17/92	1 2	26 ND	1 ND	27	0.15
	Sawtooth Weir 3/13-21/92	72,85		PAH A	CWT/LV/AD *LV/AD *AD	104422	18,336 515 152,349	5	Acclimation Experimental Group Trucked to SAWT on 3/18/92	1 2	0 ND	2 ND	2	0.01
	Sawtooth Weir 3/13-21/92	81,85		PAH A	CWT/LV/AD *LV/AD *AD	104423	19,341 810 160,699	5.1	Acclimation Experimental Group Trucked to SAWT on 3/19/92	1 2	4 ND	2 ND	6	0.03
200	Sawtooth Weir 4/10-13/92	78-80	47,895	РАН А	CWT/LV/AD *LV/AD *AD	104007	45,646 1,850 399	4.5	Size Exp Regular and Acc. Control group Trucked to & released at SAWT on 4/10/92	1 2	5 ND	4 ND	9	0.02
	Sawtooth Weir 4/10-13/92	73-77	54,645	PAH A	CWT/LV/AD *LV/AD *AD	101530	53,463 573 609	2.8	Size Exp Large Trucked to & released at SAWT on 4/10/92	1 2	265 ND	2 ND	267	0.50
	Sawtooth Weir 4/10-13/92			PAH A	NON-CWT (Includes all *	·)	467,139			1 2	162 ND	91 ND	253	0.05
	Sawtooth Weir 4/10-21/92			PAH A	**PIT		1,479		Includes all PIT tags for acc. and size exp's. See study design for complete details.	or 1 2	1 ND	0 ND	1	0.07
				TOTAL N	WT RELEASE ON-CWT RELI ITE RELEASE		154,741 467,139 621,880		TOTAL RETURN:		463	102	565	0.09

Appendix B. Table 3. Continued

					Identifying	g Marks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Return C	omposition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
TOTAL PAH A-STOC	K CWT RE	ELEASE				154,741		RETURN IS INCOM	IPLETE				
TOTAL PAH A-STOC	K NON-CV	VT RELE	ASE			690,545							
TOTAL PAH A-STOC	K RELEAS	SE				845,286	•	RETURN					
								GRAND TOTAL:		1025	241	1266	0.09
TOTAL DWOR B-STO	OCK CWT	RELEASE	Ε			55,226							
TOTAL DWOR B-STO	OCK NON-	CWT REL	EASE			547,643							
TOTAL DWOR B-STO	OCK RELE	ASE				602,869	•						
TOTAL HAGERMAN	NFH CWT	RELEAS	E			209,967							
TOTAL HAGERMAN	NFH NON-	-CWT REI	LEASE			1,238,188							
TOTAL HAGERMAN	NFH RELE	EASE				1,448,155	•						
TOTAL PIT TAGS						1,479							

Appendix C. Table 1. Release data and estimated adult returns for Magic Valley Fish Hatchery summer steelhead trout, brood year 1989. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Magic	Valley	Fish Hatch	ery Bro	od Year: 1989									
				lde	entifying N	/larks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Return C	composition	Total	SAR
Site/Date	NO.	Total	ID	Type	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Sawtooth Weir 4/12-13/90	1	1,159,080	PAH A	CWT/LV/AD/FB FB = LA-) (-1	104059	39,620	4.2	Contribution, timing	1 2	56 53	14 0	123	0.31
4/12-13/30				*LV/AD		2,804			2	00	O		
				*AD		1,156,276	4.2						
Sawtooth Weir			PAH A	NON-CWT		1,159,080		Production	1	3,312	828	5,762	0.50
4/12-13/90				(Includes all *)				(Includes all *)	2	1,424	198		
Includes all rel have Mark Typ													
, , , , , , , , , , , , , , , , , , , ,		,	TOTAL C	WT RELEASE	-	39,620							
				ON-CWT RELEAS	SE .	1,159,080		TOTAL RETURN:		4,845	1,040	5,885	0.49
			TOTAL S	ITE RELEASE		1,198,700							
E. F. Salmon R.	10	152,110	E. FK. B	CWT/LV/AD/FB	104058	40,905	4.1	Contribution, timing	1	11	4	104	0.25
4/14/90				FB = RA-)(-1					2	66	15		
				*LV/AD *AD		837 111,205			3	8	0		
E. F. Salmon R.	16	651,914	E. FK. B	CWT/LV/AD	104236	15,474	4.2	Contribution	1	13	0	37	0.24
4/16/90		•		*LV/AD		366			2	20	4		
				*AD		42,068			3	0	0		
E. F. Salmon R.	11	115,013	E. FK. B	CWT/LV/AD	104237	15,971	4.2	Contribution	1	5	1	30	0.19
4/16/90				*LV/AD		378			2	21	3		
				*AD		43,419			3	0	0		
E. F. Salmon R.	11	115,013	E. FK. B	CWT/LV/AD	104238	14,958	4.2	Contribution	1	0	3	27	0.18
4/16/90				*LV/AD		354			2	21	3		
				*AD		40,665			3	0	0		

Appendix C. Table 1. Continued.

					entifying N								
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		Composition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
E. F. Salmon R.	16	651,914	DWOR B	CWT/LV/AD	104233	14,964	5	Contribution	1	0	0	2	0.01
4/16-18/90		,-		*LV/AD		1,079			2	2	0		
				*AD		183,731			3	0	0		
E. F. Salmon R.	16	651,914	DWOR B	CWT/LV/AD	104234	15,157	5	Contribution	1	0	0	58	0.38
4/16-18/90				*LV/AD		1,093			2	58	0		
				*AD		186,101			3	0	0		
E. F. Salmon R.	16	651,914	DWOR B	CWT/LV/AD	104235	14,642	5	Contribution	1	2	0	12	0.08
4/16-18/90				*LV/AD		1,056			2	9	1		
				*AD		179,777			3	0	0		
E. F. Salmon R.			DWOR B	NON-CWT	552,837	792,129	4.6	Production	1	112	14	773	0.10
4/14-18/90			E. FK. B	(Includes all *)	239,292			(Includes all *)	2	535	73		
Includes all releas									3	39	0		
have Mark Type in	ndicate	d by *											
				VT RELEASE	-	132,071							
				N-CWT RELEAS	SE	792,129							
			TOTAL SI	TE RELEASE		924,200		TOTAL RETURN:		922	121	1,043	0.11
Slate Cr.	ND	ND	DWOR B	*AD		162,700	60.8	Excess Fingerling	1	0	0	0	0
Upper Salmon R.								Plant	2	0	0		
9/11/89					_				3	0	0		
			-	VT RELEASE	. -	0							
				ON-CWT RELEAS TE RELEASE)E	162,700 162,700		TOTAL RETURN:		0	0	0	0
			IOTAL SI	IE RELEASE		102,700		IOTAL RETURN.		U	U	U	U
TOTAL PAH A-S						39,620							
TOTAL PAH A-S			RELEASE			1,159,080							
TOTAL PAH A-S	TOCK	RELEASE				1,198,700		RETURN IS COMP	LETE				
								IXETOKITIO COM	<u></u>				
TOTAL DWOR B						44,763		DETUDN					
TOTAL DWOR B						715,537	-	RETURN		E 707	4 404	6.000	0.00
TOTAL DWOR B	-SIOC	K RELEAS	E			760,300		GRAND TOTAL:		5,767	1,161	6,928	0.30

(includes fingerling and fry releases)

Appendix C. Table 1. Continued.

				I	dentifying I	Marks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Return C	Composition	Total	SAR
Site/Date	NO.	Total	ID	Type	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
TOTAL E FK B	-ѕтоск (CWT RELE	ASE			87,308							
TOTAL E FK B	-STOCK I	NON-CWT	RELEASE			239,292							
TOTAL E FK B	-STOCK I	RELEASE				326,600	-						
TOTAL CWT R	ELEASE	FOR MAGI	C VALLEY F	Н		171,691							
TOTAL NON-C	WT RELE	ASE FOR	MAGIC VALL	EY FH		2,113,909							
TOTAL MAGIC	VALLEY	FH RELEA	SE			2,285,600	-						
TOTAL SMOLT	RELEAS	SE .				2,122,900		RETURN (includes smolt rele	eases)	5,767	1,161	6,928	0.33

Appendix C. Table 2. Release data and estimated adult returns for Magic Valley Fish Hatchery summer steelhead trout, brood year 1990. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Magic	: Valle	y Fish Hat	chery Br	ood Year: 1990									
				lo	dentifying	Marks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Return C	composition	Total	SAR
Site/Date	NO.	Total	ID	Type	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
E. F. Salmon R.	3	209,893	DWOR B	CWT/LV/AD	104314	20,498	4.1	Contribution	1	200	1	211	1.03
4/13/91				*LV/AD		2,027			2	10	0		
				*AD		187,368			3	ND	ND		
E. F. Salmon R.	2	214,470	DWOR B	CWT/LV/AD	104315	21,017	4.1	Contribution	1	5	1	70	0.33
4/13/91				*LV/AD		2,079			2	64	0		
				*AD		191,374			3	ND	ND		
E. F. Salmon R.	3	207,987	DWOR B	CWT/LV/AD	104316	20,312	4.1	Contribution	1	0	0	53	0.26
4/13/91				*LV/AD		2,008			2	53	0		
				*AD		185,667			3	ND	ND		
E. F. Salmon R.	15	113,570	E. FK. B.	CWT/LV/AD	104320	22,525	4.3	Contribution	1	4	3	24	0.11
4/15-16/91				*LV/AD		530			2	16	1		
				*AD		90,515			3	ND	ND		
E. F. Salmon R.	14	112,609	E. FK. B.	CWT/LV/AD	104321	22,483	4.3	Contribution	1	0	3	10	0.04
4/15-16/91				*LV/AD		529			2	6	1		
				*AD		89,597			3	ND	ND		
E. F. Salmon R.	15	107,771	E. FK. B.	CWT/LV/AD	104322	21,375	4.3	Contribution	1	0	1	53	0.25
4/15-16/91				*LV/AD		503			2	51	1		
				*AD		85,893			3	ND	ND		
E. F. Salmon R.	2	1,500	DWOR B	AD/PIT		1,500		Migration	1	0	0	0	0.00
								survival					
4/13, 15-16/91	14		E. FK. B.					and timing	2	0	0		
									3	ND	ND		
E. F. Salmon R.			DWOR B	NON-CWT	570,523	838,090		Production	1	337	46	1896	0.23

Includes all release dates that	E FK B	(Includes all	267,567	(Includes all *)	2	1466	47
have Mark Type indicated by * Appendix C. Table 2. Continued.		*)			3	ND	ND

				lo	lentifying	Marks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Return C	Composition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
			TOTAL N	WT RELEASE ON-CWT RELE ITE RELEASE	ASE	128,210 839,590 967,800		TOTAL RETURN:		2212	105	2317	0.24
Hazard Cr. L. Salmon R. 4/26/91	8	102,541	РАН А	CWT/LV/AD *LV/AD *AD	104317	21,809 744 79,988	3.6	Contribution	1 2	4 21	4 21	50	0.23
Hazard Cr. L. Salmon R. 4/26/91	7 8	105,146	PAH A	CWT/LV/AD *LV/AD *AD	104318	22,704 774 81,668	3.6	Contribution	1 2	21 37	21 37	116	0.51
Hazard Cr. L. Salmon R. 4/26/91	8	101,013	PAH A	CWT/LV/AD *LV/AD *AD	104319	21,484 733 78,796	3.6	Contribution	1 2	16 13	16 13	58	0.27
Hazard Cr.	7	1,600	PAH A	AD/PIT		1,600	3.6	Migration survival	1	1	1	5	0.31
L. Salmon R. 4/26/91	8							and timing	2	2	1		
Hazard Cr. L. Salmon R.			PAH A	NON-CWT (Includes all *)		242,703	3.6	Production (Includes all *)	1 2	151 261	151 261	824	0.34
Includes all relea have Mark Type				,	_								
			TOTAL N	WT RELEASE ON-CWT RELE ITE RELEASE	ASE	65,997 244,303 310,300		TOTAL RETURN:		527	526	1053	0.34
Sawtooth Weir 4/9-19/91	ND	ND	PAH A	*AD	_	364,700	3.9	Production	1 2	916 427	288 55	1686	0.46
			TOTAL N	WT RELEASE ON-CWT RELE ITE RELEASE	ASE	0 364,700 364,700		TOTAL RETURN:		1343	343	1686	0.46

Pahsimeroi R.	ND	ND	PAH A	*AD	135,100	3.8	Production	1	445	411	1159	0.86
at hatchery 4/18-19/91								2	205	98		

Appendix C. Table 2. Continued.

Appendix C. Ta			_	ld	entifying	Marks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Return C	composition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
			TOTAL NO	T RELEASE N-CWT RELEASE E RELEASE	ASE	0 135,100 135,100		TOTAL RETURN:		650	509	1159	0.86
Salmon R. at Shoup Br.	ND	ND	PAH A	*AD		97,800	3.9	Production	1	214	6	409	0.42
4/20-21/91				T RELEASE N-CWT RELEA	ASE	0 97,800			2	132	57		
			TOTAL SIT	E RELEASE		97,800		TOTAL RETURN:		346	63	409	0.42
Hammer Cr. 4/22-25/91	ND	ND	РАН А	*AD		186,300	3.9	Production	1 2	116 200	116 200	632	0.34
			TOTAL NO	T RELEASE N-CWT RELEASE E RELEASE	ASE	0 186,300 186,300		TOTAL RETURN:		316	316	632	0.34
TOTAL PAH A-S TOTAL PAH A-S TOTAL PAH A-S	STOCK	NON-CW	T RELEASE			65,997 1,028,203 1,094,200							
TOTAL DWOR I	B-STOC	K NON-C	WT RELEA	SE		61,827 571,273 633,100							
TOTAL E FK B- TOTAL E FK B- TOTAL E FK B-	STOCK	NON-CV	VT RELEASE	Ī		66,383 268,317 334,700		RETURN IS INCO	MPLETE	_			
TOTAL CWT RE TOTAL NON-CV TOTAL MAGIC	NT REL	EASE FO	R MAGIC V			194,207 1,867,793 2,062,000		GRAND TOTAL:		5,394	1,862	7,256	0.35

TOTAL PIT TAGS 3,100

Appendix C. Table 3. Release data and estimated adult returns for Magic Valley Fish Hatchery summer steelhead trout, brood year 1991. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Magic	Valle	y Fish Ha	tchery Bro	od Year: 1991	ntifying M	arke							
Release Site/Date	RW NO.	RW Total	Stock ID	Mark Type	CWT Code	Release Number	Size (FPP)	Marking Purpose	Age Ocean	Return C	Composition Hatchery	Total Returns	SAR (%)
E.F. Salmon R. 4/6-14/92	9	480,368	DWOR B	CWT/LV/AD *AD	104418	21,771 458,597	4.4	Contribution Rep. #1	1 2 3	0 ND ND	0 ND ND	0	0.00
E.F. Salmon R. 4/6-14/92	9		DWOR B	CWT/LV/AD/PIT	104418	(100)							
E.F. Salmon R. 4/6-14/92	11	476,032	DWOR B	CWT/LV/AD *LV/AD *AD	104419	21,568 143 454,321	4.5	Contribution Rep. #2	1 2 3	0 ND ND	0 ND ND	0	0.00
E.F. Salmon R. 4/6-14/92	11		DWOR B	CWT/LV/AD/PIT	104419	(100)							
E.F. Salmon R. 4/6-14/92	14	84,800	E FK B	CWT/LV/AD *AD	104420	20,821 63,979	6.1	Contribution Rep. #3	1 2 3	20 ND ND	1 ND ND	21	0.10
E.F. Salmon R. 4/6-14/92	14		E FK B	CWT/LV/AD/PIT	104420	(100)							
E.F. Salmon R. 4/6-14/92 Includes all relea have Mark Type			DWOR B E FK B	NON-CWT (Includes all *)		977,040		Production (Includes all *)	1 2 3	0 ND ND	0 ND ND	0	0.00
		0,	TOTAL NO	T RELEASE N-CWT RELEASE E RELEASE		64,160 977,040 1,041,200		TOTAL RETUR	N:	20	1	21	0.00

Appendix C. Table 3. Continued.

					ntifying M								
Release Site/Date	RW NO.	RW Total	Stock ID	Mark Type	CWT Code	Release Number	Size (FPP)	Marking Purpose	Age Ocean	Return C Harvest	Composition Hatchery	Total Returns	(%)
Hazard Cr. L. Salmon R. 4/17-25/92	4	284,715	OXBOW A	CWT/LV/AD *AD	104416	22,223 262,492	3.9	Contribution Rep. #2	1 2	5 ND	5 ND	10	0.04
Hazard Cr. L. Salmon R. 4/17-25/92	4		OXBOW A	CWT/LV/AD/PIT	104416	(100)							
Hazard Cr. L. Salmon R. 4/17-25/92	7	281,085	OXBOW A	CWT/LV/AD *AD	104417	21,604 259,481	4	Contribution Rep. #3	1 2	8 ND	8 ND	16	0.07
Hazard Cr. L. Salmon R. 4/17-25/92	7		OXBOW A	CWT/LV/AD/PIT	104417	(100)							
L. Salmon R. at Warm Springs 4/14-21/92	1 s Bridge	436100	OXBOW A	CWT/LV/AD *AD	104415	21,091 415,009	4.3	Contribution Rep. #1	1 2	5 ND	5 ND	10	0.05
L. Salmon R. at Warm Springs 4/14-21/92	1 s Bridge)	OXBOW A	CWT/LV/AD/PIT	104415	(100)							
L. Salmon R. 4/14-25/92 Includes all relea have Mark Type			OXBOW A	NON-CWT (Includes all *)		936,982	4.3	Production (Includes all *)	1 2	261 ND	261 ND	522	0.06
nave many rype	in a locat	ou 2y	TOTAL CWT	N-CWT RELEASE		64,918 936,982 1,001,900		TOTAL RETURN	N :	279	279	558	0.06
Sawtooth Weir Salmon R. 3/23-24/92	ND	ND	РАН А	*AD		117,300	5	Production	1 2	40 ND	23 ND	63	0.05
0.20 2 1102			TOTAL CW TOTAL NON TOTAL SITE	N-CWT RELEASE		0 117,300 117,300		TOTAL RETURN	N :	40	23	63	0.05

Appendix C. Table 3. Continued.

лерения с. т				lo	dentifying N	larks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Return C	Composition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
TOTAL PAH A	-ѕтоск	CWT REL	EASE			0							
TOTAL PAH A	-STOCK	NON-CW	T RELEASE			117,300		RETURN IS IN	COMPLE	TE			
TOTAL PAH A	-STOCK	RELEASE	=			117,300							
								RETURN					
TOTAL OXBO	W A-STO	OCK CWT	RELEASE			64,918		GRAND TOTA	L:	339	303	642	0.03
TOTAL OXBO	W A-STO	OCK NON-	CWT RELEASE			936,982							
TOTAL OXBO	W A-STO	OCK RELE	ASE			1,001,900							
TOTAL DWOR	B-STO	CK CWT R	ELEASE			43,339							
TOTAL DWOR	B-STO	CK NON-C	WT RELEASE			913,061							
TOTAL DWOR	B-STO	CK RELEA	SE			956,400							
TOTAL E FK B	S-STOCK	CWT RE	LEASE			20,821							
TOTAL E FK B	S-STOCK	NON-CW	T RELEASE			63,979							
TOTAL E FK B	S-STOCK	RELEAS	E			84,800							
TOTAL MAGIC	VALLE	Y FH CWT	RELEASE			129,078							
TOTAL MAGIC	VALLE	Y FH NON	-CWT RELEASE			2,031,322							
TOTAL MAGIC	VALLE	Y FH REL	EASE			2,160,400							
TOTAL PIT TA	GS					600							

Prepared by:

Approved by:

IDAHO DEPARTMENT OF FISH AND GAME

T. Dean Rhine Senior Fishery Research Biologist Virgil K. Moore, Chief Bureau of Fisheries

Randall S. Osborne Senior Fisheries Technician Steve Yundt Fishery Research Manager

Kristy A. Stevens Fisheries Biological-Aide